

# Analytical approach by Sorting of Manure for Agriland and Energy from Municipality Solid Waste

<sup>1</sup>Jitendra Kumar Srivastava,<sup>2</sup>Kamta Prasad Verma

,<sup>3</sup>Sujeet Singh,<sup>4</sup>Mannu Kumar Singh

<sup>1</sup>AP (EED) & NSS-Co-ordinator,<sup>2</sup> Lecturer (EED)

,<sup>3,4</sup> UG Scholar of EED

<sup>1,3,4</sup> B.N.College of Engineering & Technology,Lucknow,India

<sup>2</sup>Government Polytechnic,Jaunpur,India

**ABSTRACT:** Waste operation and declining soil fertility are the main issues endured by all developing nations, like India. Presently, agricultural operation of Municipal Solid Waste (MSW) is one of the most promising and cost effective options for managing solid waste. It's helpful in working current burning issues viz. soil fertility and MSW operation still there is always an implicit trouble because MSW may contain pathogens and toxic pollutants. Therefore important emphasis has been paid to composting of MSW in recent times operation of compost from MSW in agricultural land helps in upgrading the soil's physico-chemical parcels. incremental from that it also assists in perfecting natural response of cultivated land. Keeping the present situation in mind, this review critically discusses the current script, agricultural operation of MSW compost and energy from part of soil microbes and soil microbial response on external solid waste compost operation.

**Key words:** MSW, Manure, Agriland, Energy, Smart Technologies etc.

**I.INTRODUCTION:** The growing urbanization and industrialization has led to in numerous problems in developed as well as in developing countries. There are numerous pressing issues surfaced due to adding population that ultimately poses trouble to the agrarian, ecosystems and environmental sustainability either directly or laterally<sup>[2][5][7]</sup> (Fig. 1).




Fig 1: Impact of Increasing Population on Environment health

Amidst, the generation and operation of External solid waste (MSW) is important as this waste is disposed of unscientifically by low lying area without taking necessary preventives, therefore posing threat to the mortal health and near terrain. Thus there's a critical need to manage the MSW in such a way that while managing its volume and quality, it also helps to sustain the terrain<sup>[12][14]</sup>. Piecemeal from this, the environmental and health norms along with social adequacy should be achieved. Still, selection of the most applicable route for MSW operation is always being a matter of concern due to numerous environmental, specialized, fiscal, social and legislative constraints which are faced by nearly all industrially growing nations<sup>[6][11][20]</sup>.

Generally, waste generated from domestic, marketable, institutional and artificial sectors; and external services are included in MSW. MSW can be treated as renewable resource for a variety of precious products. The organic bit of MSW provides an excellent occasion for product of different value added by- products through the bio refinery conception (maximum application of waste resource)<sup>[22][25]</sup>, farther fuelling the indirect bio economy (maximizing resource effectiveness with least waste generation through which socio-profitable and environmental stability is achieved)<sup>[28][29]</sup>.

In malignancy of having numerous advantages over other conventional waste operation options, composting of MSW isn't as important as vulgarize or in the practice as it deserves in (Fig.2)<sup>[32][33]</sup>.




Fig 2: Comparison between landfill/open dumping versus composting and Sorting of MSW

This is due to lack of mindfulness and inactive programs that need to be changed. Government and original authorities should take enterprise to promote composting/ vermicomposting of organic waste<sup>[38][39]</sup>. For illustration, mindfulness juggernauts and impulses for its installation should be handed to spread this fashion at decentralized position. Also, involvement of public private cooperation (PPP) and community grounded association (CGA) should be encouraged to overcome the problem of fiscal and professionals extremity especially in the developing countries<sup>[45][47]</sup>. Piecemeal from that original authority can induce profit from better duty collection, polluters pay scheme, selling of MSW compost as being performed by Kolkata Municipal Corporation, India<sup>[55]</sup>.

## II.EXPECTED BENEFIT AND OPPORTUNITY FROM MSW COMPOST/BIOSOLIDS APPLICATION IN AGRILAND AND ELECTRICITY:

Composting of MSW has numerous advantages over inorganic diseases (IDs) whose unbridled use during last many decades has poorly affected the soil's physico-chemical and natural parcels<sup>[46][47]</sup>. Though, IDs add nutrients to the soil incontinently after operation but their long term use may change soil PH and disturb the soil microbial biota. Generally, ID tends to strain or filter down from the shops, thus requires fresh force that pollutes ground water and also emits hothouse feasts (HHFs). On contrary to this, operation of MSW compost compound shops yield and ameliorates soil nutrient profile, microbial exertion, soil texture and soften in capacity. MSW compost is rich in organic matter content, nitrogen (N) and humic substances (substantially humic acid and fulvic acid) Soil organic matter plays a significant part in maintaining soil quality, as it improves soil's physico-chemical and natural (microbial biomass) parcels. Besides this, it has high water holding capacity (WHC) and low bulk viscosity<sup>[62][65]</sup>. Humic acid in MSW compost intensifies the cation exchange capacity (CEC) and buffering capacity of soil. It has been reported by several experimenters that repeated operation of MSW compost in agrarian land helps in adding the organic matter content and C/N rate of soil in comparison to unamended soil. Therefore helps in maintaining soil fertility and its productivity. Thus, the organic toxin (like MSW compost) could be considered as a promising and sustainable volition to inorganic toxin in husbandry and horticulture<sup>[72][74]</sup>. Still, the presence of heavy essence (i.e. Cd, Cu, Zn, Pb etc.) in MSW compost are always being a matter of concern, as it can accumulate in the soil that can be absorbed by the agrarian crops which may beget variety of mortal health issues when shifted at high tropic situations through the progression of food chain. Also, in some cases these heavy essence and redundant nutrients weep through the soil and eventually pollutes beginning groundwater<sup>[66]</sup>. Increased attention of Zn, Cu and Pb in soil amended with MSW compost and set up a dwindling trend in the exertion of phosphatise and urea's conceivably due to high heavy essence attention, while dehydrogenise, catalyse and protease were remained innocent. Although, humic substances in compost act as chelating agent, therefore reduces essence solubility but it also depends on pH, swab content and cation exchange capacity (CEC) of the soil. In addition, MSW compost occasionally has high swab attention that can pose negative effect on soil texture and shops grown<sup>[67]</sup>. Others implicit pitfalls of using MSWC is presence of pathogens, and some organic compounds. Although composting is honored as a suitable treatment used for organic wastes and could inactivate several pathogens. Still, some former studies reported that some pathogens, similar as *Listeria* spp and *Salmonella* spp, have survived during the composting. Also, revealed contagious threat associated with land operation of sewage sludge in international United States and linked 43 different type of mortal contagions in sewage sludge including high cornucopia of respiratory contagions (Coronavirus, Klassevirus, and Cosavirus) with fairly lower presence of Enteroviruses. MSWC may have some organic adulterants due to the presence of ménage dangerous and artificial wastes. The presence of organic composites produced during composting of the MSW (food wastes, yard wastes and mixed paper wastes) and set up toluene, ethylbenzene, 4-dichlorobenzene, pisopropyl toluene, and naphthalene being produced in the loftiest quantities. Likewise, demonstrated presence of Polybrominated diphenyl ethers (PBDEs) in sewage sludge collected from different backwaters of Italy, which may negatively affect soil microbial biota, water cycle and mortal health when get accumulated in soil<sup>[82][83]</sup>. Piecemeal from that presence of colourful medicinals and particular care products (PPCPs) like diphenhydramine, triclosan, carbamazepine, sulfamethazine, florfenicol, levamisole, trimethoprim etc. Antibiotics like monensin, tylosin, chlortetracycline, virginiamycin, sulfamethazine is well proved in soil amended with biosolids or beast ordure. The shops have capability to accumulate PPCPs and antibiotics, therefore may pose trouble to mortal health. Effective details of MSW countries with region and application in Table 1 and Table 2 respectively.

**Table 1:**

Examples of municipal solid waste application according to the different application directions. The economic, environmental, and social impact of municipal solid waste in producing energy, electricity and fertilizer are briefly presented. Country or region refers to where the actual application was conducted for this municipal solid waste.

MSW/organic Waste Source	Soil type/pH/EC	Initial nutrient profile of Soil	Experiment type (pot/field)	Crop	Application rate	Post response of treatments	References
Valdemingomez Municipal Waste Treatment Plant Madrid, Spain	Sandy loam soil/6.4/0.1 dS m <sup>-1</sup>	OC and total N were 8.0 and 0.7 g kg <sup>-1</sup> . Similarly, P, K, Ca, Mg and Na were 0.03, 0.2, 1.5, 0.2 and 0.01 g kg <sup>-1</sup> respectively	Short term/plot	Barley	20 and 80 t ha <sup>-1</sup>	Increased microbial activity in soil; helped in maintaining long term buffering capacity of soil	Garcia-Gil et al. (2000)
Castel di Sangro, Italy	Clay/8.3	Organic C and total N were 9.7 and 1.36 g kg <sup>-1</sup> respectively	Short term/2 year field	Sugar beet and durum Wheat rotation	12 t MSW compost ha <sup>-1</sup> for sugar beet and 24 t MSW compost ha <sup>-1</sup> for durum wheat	Organic C and total N contents, dehydrogenase and nitrate reductase activities of soil increased. Dehydrogenase activity was positively correlated with b-glucosidase activity	Crecchio et al. (2001)
Municipal waste, Calcutta, India	Alluvial/5.5/0.294 d Sm <sup>-1</sup>	OC and total N were 13.9 and 1.7 g kg <sup>-1</sup> respectively	Factorial completely randomized design	N/A	0, 2.5, 10, 20 and 40 t ha <sup>-1</sup>	Substantial increase in MBC, soil respiration, urease and phosphatase activity of the soil; no adverse effect at higher dose	Bhattacharyya et al. (2003)

**Table 2:**

Examples of municipal solid waste application according to the different application directions. The economic, environmental, and social impact of municipal solid waste in producing energy, electricity, and fertilizer are briefly presented. Country or region refers to where the actual application was conducted for this municipal solid waste. Waste refers to the specific municipal solid waste used in a particular application case. The economic, environmental, and social impact refers, respectively, to cost savings performance, energy saving and emission reduction, and social behaviour gain obtained from municipal solid waste value-added applications are mentioned.

Type of waste	Country or region	Waste	Technology	Application direction	Application examples	Application impact			References
						Economic	Environmental	Social	
Municipal solid waste	Delhi, India	Biodegradable material (> 7%)	Landfill gas technology	Energy and electricity production	Collection of methane from landfills as feedstock for electricity generation	Captured methane can provide electricity for 8–18 million homes (2015)	Helping to reduce greenhouse gas emissions	Energy capture of methane to provide energy to nearby areas	(Ghosh et al. 2018)
Municipal solid waste	China	Biodegradable organic matter in municipal solid waste	Vertical gas extraction well system (low cost, most commonly used). Horizontal gas extraction well system (high efficiency, high	Energy and electricity production	Power generation and production of biogas (for vehicle and pipeline fuel) using landfill gas technology	The energy efficiency of biogas is equivalent to 228 kilotons of standard coal for energy production	Can replace 85.5% of electricity consumption or 228 kilotons of natural gas consumption (2015); can replace 90–220 kilotons of standard coal and reduce carbon dioxide emissions by	Promoting sanitary landfills as an alternative to open-air refuse collection points; producing clean energy	(Fei et al. 2019)

			cost, difficult to construct)				350–920 kilotons (2020)		
Municipal solid waste	Eskisehir, Turkey	Organic, paper, glass, plastic, metal, ash, and others	Internal combustion reciprocating engines (more broadly, high generation efficiency, and low fuel operating costs) and gas turbines technology	Energy and electricity production	Generation of electricity at landfill sites by internal combustion engines and collection of landfill gas and methane	The net present value of the maximum electrical energy value is 109,070.1 gigawatt-hour; the minimum cost of electricity generation is only \$0.054/kilowatt-hour; the price of electricity from landfill gas is \$0.133/kilowatt-hour	Has a low global warming potential (conversion of methane to carbon dioxide when used as fuel in internal combustion engines)	Promotes renewable energy development; provides waste reduction and clean energy recovery alternative	(Kale and Gökçek 2020)
Municipal solid waste	India	70% fine gravel, sand, silt, clay, 15–23% stones, brickbats, concrete fragments, 3–5% plastic, wood, textiles, and 0.9–6.5% others	Drying and sieving	Fertilizer applications	Use of soil-like material from landfills as fill (embankment, low-lying areas) and compost (horticultural, non-agricultural applications)	Reduces the need for fresh topsoil; can be used as earth fill for infrastructure projects (road and rail embankments, e.g.); low height large area fill for non-load bearing purposes (parks, golf courses); serving of lowlying areas and deep pits (mine pits)	It can be used as a low-nutrient compost in nonagricultural applications (including parks and nonfood crops); enhances nutrient growth in virgin soils	Effective reduction in old waste deposits in landfill sites (< 4.75 mm soil-like material accounts for 40–70% of total excavated waste)	(Kale and Gökçek 2020)
Municipal solid waste	Dhanbad, India	Solid organic waste	New thermal digestion technology	Fertilizer applications	Organic fertilizer production from the organic fraction of solid waste using the new thermal digestion technology	Maximum weight loss and optimum nutrient retention with minimal energy (150 °C, 135 min); total macronutrients (sodium, phosphorus, potassium) in the digested solid organic waste are above the specified standards for organic fertilizers (>1.2%)	Increased water holding capacity of organic fertilizer (43–55% increase in porosity) for plant growth; maximum nutritional value; > 90% seed germination	The effectiveness of thermal digestion in the rapid reduction in solid organic waste and nutrient cycling was demonstrated as a novel concept and research database for the clean and sustainable management of solid organic waste	(Kumar and Gupta 2021)

**III. MUNICIPAL SOLID WASTE FOR ENERGY AND ELECTRICITY PRODUCTION:** Using External solid waste for waste- to- energy pathway is necessary for waste operation and disposal. Landfill gas and anaerobic digestion are the primary styles for producing energy from external solid waste. Tip gas technology is one of the oldest and most generally used technologies for electricity generation<sup>[87][88]</sup>. The tip gas process for electricity generation comprises roughly 40 carbon dioxide and 60 methane with a high electrical and thermal energy content. In China set up a maximum tip gas value of 3.3 billion Nm<sup>3</sup> over 30 times, generating up to 7.5 billion kilowatt- hours of electricity. The minimal cost of tip gas technology for electricity generation in Turkey is only \$0.05/ kilowatt-hour.

Anaerobic digestion is able of recovering high- quality methane, converting organic waste from external solid waste into electricity and high situations of heat and working energy problems while also carrying compost and guck<sup>[91][93]</sup>. Then the waste recycling phase simplified and the tip process simplified, but it can also have a advanced power generation capacity while producing toxin and biogas as a outgrowth<sup>[99]</sup>. Estimated that using the anaerobic digestion of affordable wastes for biogas generation has the implicit to drop hothouse gas emigrations by roughly 4.36 gigatons of carbon dioxide fellow or 13 of worldwide hothouse gas emigrations from deforestation, finessed emigrations operation, crop burning, tipgas and toxin conflation emigrations. Conversion of external solid waste to energy through a waste- to- energy pathway can produce renewable energy by landing methane. For case, showed that captured methane from Delhi tips supplied 8 – 18 million houses with power in 2015. Methane traps 80 times further heat in the atmosphere than carbon dioxide does tips are releasing far more earth- warming methane into the atmosphere from the corruption of waste than preliminarily allowed a study suggests. Scientists used satellite data from four major metropolises worldwide Delhi and Mumbai in India, Lahore in Pakistan and Buenos Aires in Argentina and set up that emigrations from tips in 2018 and 2019 were 1.4 to 2.6 times advanced than earlier estimates.

When organic waste like food, wood or paper decomposes, it emits methane into the air. tips are the third- largest source of methane emigrations encyclopaedically after oil painting and gas systems and husbandry.

Although methane only accounts for about 11 of hothouse gas emigrations and lasts about a dozen times in the air, it traps 80 times further heat in the atmosphere than carbon dioxide does. Scientists estimate that at least 25 of moment's warming is driven by methane from mortal conduct.

Also, set up that a waste- to- energy factory in Taiwan, China, generated 1.33 of original electricity consumption, with anticipated electricity product effectiveness of 30, corresponding to 346 and 748 gigawatt- hours by 2030 and 2060, independently. likewise, suggested combined cogeneration of hydrogen from electrolysis and power from the anaerobic digestion process<sup>[102][105]</sup>.

In addition to renewable energy- to- energy generation has the implicit to reduce hothouse gas emissions. The environmental performance of mongrel and tip gas blending styles in the Nigerian region, with hothouse emigration reduction rates of 76 – 93 and 75 – 85, independently. In addition, using the waste- to- energy conception can save on reactionary energy combustion and significantly reduce the cost of electricity generation. The minimal price of electricity generation is only \$0.054/kilowatt- hour compared to \$0.133/kilowatt- hour for tipgas, with a significant reduction in the total quantum of inclined waste. In addition, the waste- to- energy conception provides a way to reclaim, exercise and add value to waste, provides an volition to clean energy recovery and facilitates the sustainable development of druthers to reactionary energy combustion. Both anaerobic digestion and tip gas technologies have good environmental, profitable and social performance for electricity generation. Still, showed that anaerobic digestion has a advanced and further profitable eventuality for electricity generation than tip gas in the study area<sup>[107][109]</sup>. Delved the power generation eventuality of tip gas and anaerobic digestion technologies in Tehran and Beijing over 20 times. They set up that the technologies generated 45.2 and 41.9 further electricity than tip gas technologies in Tehran and Beijing independently and that anaerobic digestion had the most substantial eventuality to alleviate global warming. Therefore anaerobic digestion has tremendous eventuality for producing power from external solid waste. Tip waste treatment styles presently face the challenge of inefficiency, particularly when treating food waste comprising over 60 of the water content. One approach to working this issue is by reducing the food waste content of waste incineration; for illustration, reducing the waste's water content by 9 – 44 significantly increased spicy value and thus bettered power generation effectiveness.

In conclusion, using waste- to- energy is the stylish way to dispose of and add value to waste to meet the growing world population and the adding volume of external solid waste. At the same time the product of clean renewable energy as an volition to fossil energies creates a righteous cycle in profitable, environmental and social terms contributing to the development of sustainable metropolises and a global green future<sup>[111][112]</sup>.

**IV. MUNICIPAL SOLID WASTE FOR FERTILIZER APPLICATION:** Uses of inorganic nitrogen comprise about 50 of current agrarian product; still, the heavy use of inorganic diseases poses climate and environmental enterprises. For illustration, inorganic diseases contribute to large quantities of hothouse gas emigrations and eutrophication of the water terrain<sup>[116]</sup>. On the other hand, organic diseases can ameliorate organic carbon in the soil while furnishing sufficient nutrients to shops. Thus replacing inorganic diseases with organic diseases is urgently demanded to address current environmental issues.

External solid waste can be used either to produce high- quality liquid diseases from organic waste or excerpt soil- such like accoutrements from organic waste for tip and toxin use. Several recent studies have shown the possibilities of producing organic diseases from external waste<sup>[126][142]</sup>. For illustration, proposed the birth of organic carbon from external solid waste compost technology to produce 200 L of liquid toxin at €1/litre per 100 kg of dry compost verified the possibility of rooting organic substances from external solid waste's organic bit and accelerating methane product. Birth technologies of high- value organic toxin from external solid waste are entered further attention and invention at a lower cost Conventional detergent and microwave oven- supported birth are common for liquid diseases<sup>[146][148]</sup>. The birth of liquid diseases by alkaline traditional solvent birth ways is a simple, effective and ecofriendly system. In addition, traditional solvent birth is a lower energy- needed ferocious system, with a selling cost of €1/ litre and the toxin yield is ten times advanced than that of water- grounded birth.

Microwave oven- supported birth is considered a more environmental friendly and green technology than conventional solvent birth. Still, microwave oven- supported birth requires more complex conditions during the birth process, similar as advanced temperatures, power and limitations in the dielectric parcels of solid accoutrements. Microwave oven- supported birth is similar to conventional solvent birth ways when adding the operating temperature and reducing the response time<sup>[153]</sup>.

In general, the liquid diseases produced from external solid waste have much advanced total macronutrients( sodium, phosphorus, potassium) than those specified for organic diseases, ameliorate soil water- holding capacity, increase porosity and benefit factory of crop growth<sup>[163]</sup>. The new thermal digestion is a new type of digestion that has been developed to make the operation of organic

diseases from the organic bit of solid waste more effective and environmentally friendly, hence achieving maximum weight loss of waste and optimum nutrient retention of toxin with minimal energy consumption within 135 twinkles at 150 °C.

In addition, soil- suchlike material from external solid waste piles can be used as filler for road dikes and low- lying areas, Compost for horticulture and other non-agricultural operations. Through the relinquishment of this technology, the total quantum of waste in tips is significantly reduced, reducing the need for fresh soil and saving on tip costs and waste operation and disposal costs<sup>[173][181]</sup>. Considering the possible presence of heavy essence ions in soil- suchlike accoutrements in waste piles, their use for non-edible crops can reduce their threat and hazard while enhancing the nutrient content of virgin soil for non-agricultural operations. Although the feasibility of organic birth from the external solid waste operation has been vindicated, the technology's trustability and the liquid toxin quality still need to be supported by a lot of exploration data<sup>[187][189]</sup>. In addition, applying external solid waste to prize organic liquid diseases still needs important disquisition. Using other organic remainders as raw accoutrements in cutting edge technology.

In conclusion, using more advanced technologies to prize high- quality liquid diseases from the organic bit of external solid waste and using soil- suchlike accoutrements from external solid waste as compost for tip and non-agricultural operations are excellent styles for the valorisation of external solid waste<sup>[176]</sup>. Such an approach in the direction of toxin operations provides a new conception, innovative technology, and a dependable pool of exemplifications for the clean and sustainable operation of solid organic waste.

This section explains the rearmost directions in applying external solid waste in energy, electricity product and toxin and demonstrates system feasibility<sup>[90]</sup>. The exercise of external solid waste isn't only outstanding for generating electricity from waste but also for the significant mitigation of the hothouse effect and the product and negotiation of new energy sources at a lower cost. In addition, external solid waste also performs well in the medication of liquid diseases. Technological inventions have been applied to achieve minimum energy consumption to achieve maximum waste consumption and optimum nutrient retention, reduce product costs and increase the effectiveness of toxin product likewise, treated waste in tip reduces the total quantum of waste, reduce the use of fresh soil and ameliorate soil nutrients. It offers innovative results for clean energy recovery and renewable energy development operations, furnishing the rearmost technology and indefatigable power for value- adding and operation of external solid waste<sup>[138]</sup>.

**V.SOLID WASTE PRETREATMENT:** Recycling and sorting is the first and critical step in the valorisation and operation of waste programs and installations should ameliorate waste's recovery rate and sorting delicacy. First, the policy section on waste recycling and sorting should be as detailed as possible, down to the unit responsible for enforcing the policy and the rules and regulations<sup>[64][91][136]</sup>. The approach should also suit the characteristics of the region where waste is enforced. Second, waste recycling and sorting installations should also consider the drivers' age and height to make the installations universal simple and effective. In sitting installations, spatial analysis of geographic information systems can be used to screen and identify the most suitable areas or locales for recovering installations. At the same time, governments non-governmental and other associations should concentrate on changing consumer waste in the future<sup>[138]</sup>.

Publicize the negative profitable, environmental and social impacts of magpie waste disposal and call for and companion consumers to reclaim and separate their waste effectively. To achieve early results, consideration could also be given to adding the demand to reclaim and separate waste to the citizens' law of conduct to raise mindfulness of citizens' power<sup>[104]</sup>. In addition the government can also encourage businesses to develop recycling programs for vended goods linked to consumers' waste recycling geste. Also a detailed bracket and characterization of a specific type of artificial solid waste could be conducted. The suggestion of green desulfurization of scrap tires is in line with the indirect frugality and the product of rubber- grounded accoutrements for high- value end ground tires requests will be developed because of current exploration trends. Still, proper sorting and acceptable characterization of scrap rubber before use can significantly ameliorate the process reproducibility and the performance parcels of the attained rubber recycling products<sup>[152]</sup>. In addition to this, suggested that maybe in the future, the perpetration of independent robotic systems for waste recycling could be achieved with automatic sorting and physical sorting of recyclables according to material type. However, this will significantly ameliorate effectiveness and delicacy of recycling and prepare the waste for exercise, If artificial intelligence can be successfully spread to the waste recycling field<sup>[81]</sup>.

In conclusion, recovering sorting technology can ameliorate the recycling rate of waste. The help of a policy system effectively facilitates the recycling of waste. Also, recovering according to the nature of specific waste will increase the delicacy of waste recycling.



Fig 3: Enhanced recycling and sorting techniques for solid waste contribute to more efficient waste applications

This section summarizes the prospects for value- added solid waste operations, as shown in Fig.3. This graph illustrates the solid waste value- added openings in terms of operations, profitable feasibility assessment styles and the sorting direction of solid waste recycling. It's determined how sorting technology for waste recycling can be bettered. There's also a need to expand solid waste operations with added value. Some new evaluation styles and profitable parameters can be added to increase the chances of profitable viability.

**VI. PROPOSED PROTOTYPE CONCEPT:** In this MSW sorted manure used for agriland and prepared fuel used for furnace fuel. Furnace heat the boiler then steam generated. Steam pressure comes on the turbine after this turbine rotate and produces mechanical energy that connected by belt system with dynamo. Dynamo produces electricity then transmitted to grid.



Fig 4: A prototype structure for generation of electricity by MSW Based Fuel

**VII. RESULTS:** The conversion of external waste into value-added products holds excellent pledge for Agriland and Electricity. However, applicable technologies for effective conversion are still lacking and the specialized walls are substantially due to the diversity of the waste. Unborn experimenters may need to borrow indispensable exploration styles to circumvent the unreliable goods of waste diversity<sup>[217]</sup>. Solid waste exercise in different operation directions, technological invention and unborn sustainable development and furnishing a library of styles for the profitable assessment of actors in the field of solid waste<sup>[138]</sup>. Pyrolysis units for putrefying external solid waste are precious and bear a lot of thermal energy. The integration into the pyrolysis unit can minimize this pyrolysis heating problem and make the system more eco-friendly and energy effective.

**VIII. CONCLUSION:** While accelerated global urbanization, technological inventions in husbandry and the expansion of artificial robotization have contributed to mortal development and progress, they've brought more solid waste, accelerating the environmental extremity and energy problems. This review analyzes and summarizes economically feasible styles for valorising solid waste from external, agrarian and artificial sources grounded on the rearmost reusing and value-added technologies. Tips are no longer the primary system of solid waste disposal; new ways of solid waste disposal have set up a way to misbehave with sustainable green development. For case, using solid waste as an indispensable energy source for power generation is one of the most common ways of dealing with solid waste, achieving a positive impact on global warming. In addition, solid wastes can be used for toxin operations, factory parentage, construction material- oil painting, biomethane for machine energy, biochar for soil remediation, biosorbents for wastewater treatment, beast feed, accoutrements for water storehouse systems and conservative natural coffers. As per proposed method manure development for farming as well as energy. Therefore, energy or by products can be attained at a lower cost to maximize solid waste application and cover mortal health the terrain and natural coffers. More importantly, combining value assessment and profitable feasibility analysis is vital to optimizing the profitable benefits of solid waste exercise in different operation directions, technological invention and unborn sustainable development and furnishing a library of styles for the profitable assessment of actors in the field of solid waste.

Despite the significance of recovering waste to realize value, current programs and installations for recycling and application of waste aren't well advanced and there are significant limitations in the measures taken to exercise solid waste in several countries. Therefore, there will be further room for advancement in the future in the disquisition of operations and technological invention in solid waste recycling to maximize the value added and application of solid waste.

## IX. REFERENCES:

- [1] Abdollahi Saadatlu E et al (2022) A sustainable model for municipal solid waste system considering global warming potential impact: a case study. *Comput Ind Eng* 169:108127. <https://doi.org/10.1016/j.cie.2022.108127>
- [2] Abdullah I et al (2022) Conversion of biomass blends (walnut shell and pearl millet) for the production of solid biofuel via torrefaction under different conditions. *Chemosphere* 295:133894. <https://doi.org/10.1016/j.chemosphere.2022.133894>
- [3] Abdulyekeen KA et al (2021) Torrefaction of biomass: production of enhanced solid biofuel from municipal solid waste and other types of biomass. *Renew Sustain Energy Rev* 150:111436. <https://doi.org/10.1016/j.rser.2021.111436>
- [4] Abedinzadeh M et al (2020) Combined use of municipal solid waste biochar and bacterial biosorbent synergistically decreases Cd (II) and Pb (II) concentration in edible tissue of forage maize irrigated with heavy metal-spiked water. *Heliyon* 6:e04688. <https://doi.org/10.1016/j.heliyon.2020.e04688>
- [5] Abraham JJ et al (2022) An experimental study on concrete block using construction demolition waste and life cycle cost analysis. *Mater Today: Proc* 60:1320–1324. <https://doi.org/10.1016/j.matpr.2021.09.307>
- [6] Afroze S et al (2018) A review on heavy metal ions and dye adsorption from water by agricultural solid waste adsorbents. 229,1-50. <https://doi.org/10.1007/s11270-018-3869-z>
- [7] Ahmad T et al (2019) Treatment and utilization of dairy industrial waste: a review. *Trends Food Sci Technol* 88:361–372. <https://doi.org/10.1016/j.tifs.2019.04.003>
- [8] Ahmed MJK, Ahmaruzzaman M (2016) A review on potential usage of industrial waste materials for binding heavy metal ions from aqueous solutions. *J Water Process Eng* 10:39–47. <https://doi.org/10.1016/j.jwpe.2016.01.014>
- [9] Akinrinmade AO (2020) Determination of appropriate landfill sites and materials in parts of Kwara State Nigeria using geological, geophysical and geotechnical techniques. Kwara State University (Nigeria)
- [10] Akor CI et al (2021) Thermokinetic study of residual solid digestate from anaerobic digestion. *Chem Eng J* 406:127039. <https://doi.org/10.1016/j.cej.2020.127039>

- [11]Al-Anzi FS (2022) Building a planter system using waste materials using value engineering environmental assessment. *Sci Rep* 12:2344. <https://doi.org/10.1038/s41598-022-05300-0>
- [12]Almendro-Candel MB et al (2018) Physical properties of soils affected by the use of agricultural waste. <https://doi.org/10.5772/intech.hopen.77993>
- [13]Al-Osta MA et al (2016) Study of heavy fuel oil fly ash for use in concrete blocks and asphalt concrete mixes. *Adv Concr Constr* 4:123. <https://doi.org/10.12989/acc.2016.4.2.123>
- [14]Anand S (2010) Solid waste management. Mittal Publications. Anastopoulos I et al (2019) Valorization of agricultural wastes could improve soil fertility and mitigate soil direct N<sub>2</sub>O emissions. *J Environ Manage* 250:109389. <https://doi.org/10.1016/j.jenvman.2019.109389>
- [15]Arafat HA et al (2015) Environmental performance and energy recovery potential of five processes for municipal solid waste treatment. *J Clean Pro* 105:233–240. <https://doi.org/10.1016/j.jclepro.2013.11.071>
- [16]Arpia AA et al (2021) Sustainable biofuel and bioenergy production from biomass waste residues using microwave-assisted heating: a comprehensive review. *Chem Eng J* 403:126233. <https://doi.org/10.1016/j.cej.2020.126233>
- [17]Asif M et al (2017) Potential of chitosan alone and in combination with agricultural wastes against the root-knot nematode, *Meloidogyne incognita* infesting eggplant. *J Plant Prot Res* 57:288–295. <https://doi.org/10.1515/jppr-2017-0041>
- [18]Awasthi MK et al (2021a) A critical review on the development stage of biorefinery systems towards the management of apple processing-derived waste. *Renew Sustain Energy Rev* 143:110972. <https://doi.org/10.1016/j.rser.2021.110972>
- [19]Awasthi MK et al (2021b) Techno-economics and life-cycle assessment of biological and thermochemical treatment of bio-waste. *Renew Sustain Energy Rev* 144:110837. <https://doi.org/10.1016/j.rser.2021.110837>
- [20]Ayeleru OO et al (2021) Cost benefit analysis of a municipal solid waste recycling facility in Soweto. *South Africa Waste Manag* 134:263–269. <https://doi.org/10.1016/j.wasman.2021.08.001>
- [21]Ayodele T et al (2017) Life cycle assessment of waste-to-energy (WtE) technologies for electricity generation using municipal solid waste in Nigeria. *Appl Energy* 201:200–218. <https://doi.org/10.1016/j.apenergy.2017.05.097>
- [22]Azam M et al (2019) Status, characterization, and potential utilization of municipal solid waste as renewable energy source: Lahore case study in Pakistan. *Environ Int* 134:105291. <https://doi.org/10.1016/j.envint.2019.105291>
- [23]Banerjee M et al (2019) Cu (II) removal using green adsorbents: kinetic modelling and plant scale-up design. *Environ Sci Pollut Res* 26:11542–11557. <https://doi.org/10.1007/s11356-018-1930-5>
- [24]Batista Meneses D et al (2022) Pretreatment methods of lignocellulosic wastes into value-added products: recent advances and possibilities. *Biomass Convers Biorefinery* 1:547–564. <https://doi.org/10.1007/s13399-020-00722-0>
- [25]Bayık GD, Altın A (2018) Conversion of an industrial waste to an oil sorbent by coupling with functional silanes. *J Clean Prod* 196:1052–1064. <https://doi.org/10.1016/j.jclepro.2018.06.076>
- [26]Bekchanov M, Mirzabaev A (2018) Circular economy of composting in Sri Lanka: opportunities and challenges for reducing waste related pollution and improving soil health. *202:1107–1119*. <https://doi.org/10.1016/j.jclepro.2018.08.186>
- [27]Bernal V et al (2018) Physicochemical properties of activated carbon: their effect on the adsorption of pharmaceutical compounds and adsorbate-adsorbent interactions. *C-J Carbon Res* 4:62. <https://doi.org/10.3390/c4040062>
- [28]Bernat K et al (2022) Can the biological stage of a mechanical-biological treatment plant that is designed for mixed municipal solid waste be successfully utilized for effective composting of selectively collected biowaste? *Waste Manage* 149:291–301. <https://doi.org/10.1016/j.wasman.2022.06.025>
- [29]Bharathiraja B et al (2018) Biogas production—a review on composition, fuel properties, feed stock and principles of anaerobic digestion. *Renew Sustain Energy Rev* 90:570–582. <https://doi.org/10.1016/j.rser.2018.03.093>
- [30]Bhattacharya A et al (2021) Taxonomy of antecedents of food waste—a literature review. *J Clean Prod* 291:125910. <https://doi.org/10.1016/j.jclepro.2021.125910>
- [31]Bieliatynskiy A et al (2022) The use of fiber made from fly ash from power plants in China in road and air field construction. *Constr Build Mater* 323:126537. <https://doi.org/10.1016/j.conbuildmat.2022.126537>
- [32]Bisaglia C et al (2018) Methane/Gasoline Bi-fuel engines as a power source for standard agriculture tractors: development and testing activities. *Appl Eng Agric* 34:365–375. <https://doi.org/10.13031/aea.12262>
- [33]Breunig H et al (2022) Economic and greenhouse gas analysis of regional bioenergy-powered district energy systems in California. *Resour Conserv Recy* 180:106187. <https://doi.org/10.1016/j.resconrec.2022.106187>
- [34]Brigde J, Starr J (2007) Plant nematodes of agricultural importance. Manson publishing Ltd. P., <https://doi.org/10.1201/b15142>
- [35]Cai J et al (2021) Coupling and coordinated development of new urbanization and agro-ecological environment in China. *Sci Total Environ* 776:145837. <https://doi.org/10.1016/j.scitotenv.2021.145837>
- [36]Caiardi F et al (2022) Waste-to-energy innovative system: assessment of integrating anaerobic digestion and pyrolysis technologies. *Sustain Prod Consum* 31:657–669. <https://doi.org/10.1016/j.spc.2022.03.021>
- [37]Campos EV et al (2019) Use of botanical insecticides for sustainable agriculture: future perspectives. *Ecol Ind* 105:483–495. <https://doi.org/10.1016/j.ecolind.2018.04.038>
- [38]Campuzano R, González-Martínez S (2017) Influence of process parameters on the extraction of soluble substances from OFMSW and methane production. *Waste Manage* 62:61–68. <https://doi.org/10.1016/j.wasman.2017.02.015>
- [39]Cao Y et al (2022) Development of a MSW-fueled sustainable co-generation of hydrogen and electricity plant for a better environment comparing PEM and alkaline electrolyzers. *Sustain Cities Soc* 81:103801. <https://doi.org/10.1016/j.scs.2022.103801>
- [40]Cetrulo TB et al (2018) Effectiveness of solid waste policies in developing countries: a case study in Brazil. *J Clean Prod* 205:179–187. <https://doi.org/10.1016/j.jclepro.2018.09.094>
- [41]Chaianong A, Pharino C (2022) How to design an area-based prioritization of biogas production from organic municipal solid waste? *Evid Thailand Waste Manag* 138:243–252. <https://doi.org/10.1016/j.wasman.2021.11.042>
- [42]Chehade G, Dincer I (2021) Progress in green ammonia production as potential carbon-free fuel. *Fuel* 299:120845. <https://doi.org/10.1016/j.fuel.2021.120845>
- [43]Chen Y-C, Lo S-L (2016) Evaluation of greenhouse gas emissions for several municipal solid waste management strategies. *J Clean Prod* 113:606–612. <https://doi.org/10.1016/j.jclepro.2015.11.058>



- [44]Chen Z et al (2010) Overview on LFG projects in China. *Waste Manage* 30:1006–1010. <https://doi.org/10.1016/j.wasman.2010.02.001>
- [45]Chen T et al (2015) A safety analysis of food waste-derived animal feeds from three typical conversion techniques in China. *Waste Manage* 45:42–50. <https://doi.org/10.1016/j.wasman.2015.06.041>
- [46]Chen L et al (2022) Strategies to achieve a carbon neutral society:a review. *Environ Chem Lett* 20:2277–2310. <https://doi.org/10.1007/s10311-022-01499-6>
- [47]Commission E (2012) Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. *Innovating for sustainable growth: a bioeconomy for Europe.*, Brussels, Belgium.
- [48]Crini G, Lichtfouse E (2018) Advantages and disadvantages of techniques used for wastewater treatment. *Environ Chem Lett* 17:145–155. <https://doi.org/10.1007/s10311-018-0785-9>
- [49]Cudjoe D et al (2020) Electricity generation using biogas from organic fraction of municipal solid waste generated in provinces of China: techno-economic and environmental impact analysis. *Fuel Process Technol* 203:106381. <https://doi.org/10.1016/j.fuproc.2020.106381>
- [50]Cudjoe D et al (2021a) Power generation from municipal solid waste landfilled in the Beijing-Tianjin-Hebei region. *Energy* 217:119393. <https://doi.org/10.1016/j.energy.2020.119393>
- [51]Cudjoe D et al (2021b) Power generation from municipal solid waste landfilled in the Beijing-Tianjin-Hebei region. *Energy* 217:119393. <https://doi.org/10.1016/j.energy.2020.119393>
- [52]Dai Y et al (2018) Utilizations of agricultural waste as adsorbent for the removal of contaminants: a review. *Chemosphere* 211:235–253. <https://doi.org/10.1016/j.chemosphere.2018.06.179>
- [53]Dao TAT et al (2020) Optimization of pectin extraction from fruit peels by response surface method: conventional versus microwave-assisted heating. *Food Hydrocoll* 113:106475. <https://doi.org/10.1016/j.foodhyd.2020.106475>
- [54]Das S et al (2019) Solid waste management: Scope and the challenge of sustainability. *J Clean Prod* 228:658–678. <https://doi.org/10.1016/j.jclepro.2019.04.323>
- [55]Datta M et al (2021) Feasibility of re-using soil-like material obtained from mining of old MSW dumps as an earth-fill and as compost. *Process Saf Environ Prot* 147:477–487. <https://doi.org/10.1016/j.psep.2020.09.051>
- [56]De Gisi S et al (2016) Characteristics and adsorption capacities of low-cost sorbents for wastewater treatment: a review. *Sustain Mater Technol* 9:10–40. <https://doi.org/10.1016/j.susmat.2016.06.002>
- [57]Demirbas A (2007) Bio-fuels from agricultural residues. *Energy Sources, Part a: Recovery, Utilization, Environ Ef* 30:101–109. <https://doi.org/10.1080/00908310600626788>
- [58]Deniz F, Kepekci RA (2016). Dye biosorption onto pistachio byproduct: a green environmental engineering approach. *J Mol Liq* 219:194–200. <https://doi.org/10.1016/j.molliq.2016.03.018>
- [59]Di Maria F et al (2021) The life cycle approach for assessing the impact of municipal solid waste incineration on the environment and on human health. *Sci Total Environ* 776:145785. <https://doi.org/10.1016/j.scitotenv.2021.145785>
- [60]Diaz LF et al 2011. *Compost science and technology*. Elsevier Dlamini S et al (2019) Municipal solid waste management in South Africa: from waste to energy recovery through waste-to-energy technologies in Johannesburg. *Local Environ* 24:249–257. <https://doi.org/10.1080/13549839.2018.1561656>
- [61]Dorward LJ (2012) Where are the best opportunities for reducing greenhouse gas emissions in the food system (including the food chain)? *Food policy*. 37:463–466. <https://doi.org/10.1016/j.foodpol.2012.04.006>
- [62]Duan Y et al (2020) Organic solid waste biorefinery: sustainable strategy for emerging circular bioeconomy in China. *Ind Crop Prod* 153:112568. <https://doi.org/10.1016/j.indcrop.2020.112568>
- [63]Dumlao-Tan MI, Halog A (2017) Moving towards a circular economy in solid waste management: concepts and practices. *Adv Solid Hazard Waste Manag*. 29–48. [https://doi.org/10.1007/978-3-319-57076-1\\_2](https://doi.org/10.1007/978-3-319-57076-1_2)
- [64]Duque-Acevedo M et al (2020) Agricultural waste: review of the evolution, approaches and perspectives on alternative uses. *Global Ecol Conserv* 22:e00902. <https://doi.org/10.1016/j.gecco.2020.e00902>
- [65]El-Azazy M et al (2019) Potato peels as an adsorbent for heavy metals from aqueous solutions: eco-structuring of a green adsorbent operating Plackett-Burman design. *J Chem*. <https://doi.org/10.1155/2019/4926240>
- [66]Elnagmy A et al (2018) Biogeological radiation hazards of some fertilizer brands in upper Egypt. *Assiut Univ J Multidiscip Sci Res* 47:21–40. <https://doi.org/10.21608/aunj.2018.221229>
- [67]Enaime G et al (2020) Biochar for wastewater treatment—conversion technologies and applications. *Appl Sci* 10:3492. <https://doi.org/10.3390/app10103492>
- [68]Eriksson M et al (2015) Carbon footprint of food waste management options in the waste hierarchy—a Swedish case study. *J Clean Pro* 93:115–125. <https://doi.org/10.1016/j.jclepro.2015.01.026>
- [69]Fabiya OA et al (2018) Suppression of heterodera sacchari in rice with agricultural waste-silver nano particles. *J Solid Waste Technol Manag* 44:87–91. <https://doi.org/10.5276/JSWTM.2018.87>
- [70]Farghali M et al (2022) Seaweed for climate mitigation, wastewater treatment, bioenergy, bioplastic, biochar, food, pharmaceuticals and cosmetics: a review. *Environ Chem Lett*. <https://doi.org/10.1007/s10311-022-01520-y>
- [71]Fausto-Castro L et al (2020) Selection of food waste with low moisture and high protein content from Mexican restaurants as a supplement to swine feed. *J Clean Pro* 256:120137. <https://doi.org/10.1016/j.jclepro.2020.120137>
- [72]Fei F et al (2019) Spatio-temporal estimation of landfill gas energy potential: a case study in China. *Renew Sustain Energy Rev* 103:217–226. <https://doi.org/10.1016/j.rser.2018.12.036>
- [73]Fernández-Delgado M et al (2020) Recovery of organic carbon from municipal mixed waste compost for the production of fertilizers. *J Clean Prod* 265:121805. <https://doi.org/10.1016/j.jclepro.2020.121805>
- [74]Fernández-Delgado M et al (2022) Liquid fertilizer production from organic waste by conventional and microwave-assisted extraction technologies: techno-economic and environmental assessment. *Sci Total Environ* 806:150904. <https://doi.org/10.1016/j.scitoenv.2021.150904>

- [75] Ferronato N et al (2021) Sensitivity analysis and improvements of the recycling rate in municipal solid waste life cycle assessment: focus on a Latin American developing context. *Waste Manage* 128:1–15. <https://doi.org/10.1016/j.wasman.2021.04.043>
- [76] Filimonau V et al (2022) Exploring the potential of industrial symbiosis to recover food waste from the foodservice sector in Russia. *Sustainable Production and Consumption* 29:467–478. <https://doi.org/10.1016/j.spc.2021.10.028>
- [77] Forghani F, Hajihassani A (2020) Recent advances in the development of environmentally benign treatments to control root-knot nematodes. *Front Plant Sci* 11:1125. <https://doi.org/10.3389/fpls.2020.01125>
- [78] Gao M et al (2019) Biogas potential, utilization and countermeasures in agricultural provinces: a case study of biogas development in Henan Province China. *Renew Sustain Energy Rev* 99:191–200. <https://doi.org/10.1016/j.rser.2018.10.005>
- [79] Georganas A et al (2020) Bioactive compounds in food waste: a review on the transformation of food waste to animal feed. *Foods* 9:291. <https://doi.org/10.3390/foods9030291>
- [80] Ghosh P et al (2018) Assessment of methane emissions and energy recovery potential from the municipal solid waste landfills of Delhi, India. *Bioresour Technol* 272:611–615. <https://doi.org/10.1016/j.biortech.2018.10.069>
- [81] Gil A (2022) Challenges on waste-to-energy for the valorisation of industrial wastes: electricity, heat and cold, bioliquids and biofuels. *Monitoring, Management* 17:100615. <https://doi.org/10.1016/j.enmm.2021.100615>
- [82] Godinho D et al (2019) Recovery of Cr (III) by using chars from the co-gasification of agriculture and forestry wastes. *Environ Sci Pollut Res* 26:22723–22735. <https://doi.org/10.1007/s11356-019-05609-w>
- [83] Gonzalez PGA et al (2022) Soybean straw as a feedstock for value added chemicals and materials: recent trends and emerging prospects. *BioEnergy Res*. <https://doi.org/10.1007/s12155-022-10506-1>
- [84] Gopalakrishnan PK et al (2021) Cost analysis and optimization of Blockchain-based solid waste management traceability system. *Waste Manage* 120:594–607. <https://doi.org/10.1016/j.wasman.2020.10.027>
- [85] Gopikumar S et al (2021) A method of landfill leachate management using internet of things for sustainable smart city development. *Sustain Cities Soc* 66:102521. <https://doi.org/10.1016/j.scs.2020.102521>
- [86] Gravert TKO et al (2021) Non-target analysis of organic waste amended agricultural soils: characterization of added organic pollution. *Chemosphere* 280:130582. <https://doi.org/10.1016/j.chemosphere.2021.130582>
- [87] Guan Y et al (2019) Dynamic analysis of industrial solid waste metabolism at aggregated and disaggregated levels. *J Clean Prod* 221:817–827. <https://doi.org/10.1016/j.jclepro.2019.01.271>
- [88] Guedes RE et al (2018) Operating parameters for bio-oil production in biomass pyrolysis: a review. *J Anal Appl Pyrol* 129:134–149. <https://doi.org/10.1016/j.jaap.2017.11.019>
- [89] Gujre N et al (2021) Deciphering the dynamics of glomalin and heavy metals in soils contaminated with hazardous municipal solid wastes. *J Hazard Mater* 416:125869. <https://doi.org/10.1016/j.jhazmat.2021.125869>
- [90] Guo J et al (2021) Improving benzo pyrene biodegradation in soil with wheat straw-derived biochar amendment: performance, microbial quantity, CO<sub>2</sub> emission, and soil properties. *J Anal Appl Pyrol* 156:105132. <https://doi.org/10.1016/j.jaap.2021.105132>
- [91] Gwenzi W et al (2015) Biochar production and applications in sub Saharan Africa: opportunities, constraints, risks and uncertainties. *J Environ Manage* 150:250–261. <https://doi.org/10.1016/j.jenvman.2014.11.027>
- [92] Haile A et al (2021) Pulp and paper mill wastes: utilizations and prospects for high value-added biomaterials. *Bioresour Bioprocess* 8:1–22. <https://doi.org/10.1186/s40643-021-00385-3>
- [93] Haqq-Misra J et al (2022) Future of life in the solar system and beyond. *New Frontiers in Astrobiology* 255–283. <https://doi.org/10.1016/B978-0-12-824162-2.00001-4>
- [94] Hasan MM et al (2021) Energy recovery from municipal solid waste using pyrolysis technology: a review on current status and developments. *Renew Sustain Energy Rev* 145:111073. <https://doi.org/10.1016/j.rser.2021.111073>
- [95] He X (2017) Information on impacts of climate change and adaptation in China. *J Environ Inf*. <https://doi.org/10.3808/jei.201700367>
- [96] Helliwell R, Burton RJF (2021) The promised land? Exploring the future visions and narrative silences of cellular agriculture in news and industry media. *J Rural Stud* 84:180–191. <https://doi.org/10.1016/j.jrurstud.2021.04.002>
- [97] Huang W, Fooladi H (2021) Economic and environmental estimated assessment of power production from municipal solid waste using anaerobic digestion and landfill gas technologies. *Energy Rep* 7:4460–4469. <https://doi.org/10.1016/j.egy.2021.07.036>
- [98] Ighalo JO et al (2022) Flash pyrolysis of biomass: a review of recent advances. *Clean Technology Environment Policy*. <https://doi.org/10.1007/s10098-022-02339-5>
- [99] Jabeen F et al (2022) Trash to energy: a measure for the energy potential of combustible content of domestic solid waste generated from an industrialized city of Pakistan. *J Taiwan Inst Chem E* 137:104223. <https://doi.org/10.1016/j.jtice.2022.104223>
- [100] Janyasuthiwong S et al (2015) Copper, lead and zinc removal from metal-contaminated wastewater by adsorption onto agricultural wastes. *Environment Technology* 36:3071–3083. <https://doi.org/10.1080/09593330.2015.1053537>
- [101] Kaab A et al (2019) Combined life cycle assessment and artificial intelligence for prediction of output energy and environmental impacts of sugarcane production. *Sci Total Environ* 664:1005–1019. <https://doi.org/10.1016/j.scitotenv.2019.02.004>
- [102] Kadier A et al (2021) Use of industrial wastes as sustainable nutrient sources for bacterial cellulose (BC) production: mechanism, advances, and future perspectives. *Polymers* 13:3365. <https://doi.org/10.3390/polym13193365>
- [103] Kainthola J et al (2019) A review on enhanced biogas production from anaerobic digestion of lignocellulosic biomass by different enhancement techniques. *Process Biochem* 84:81–90. <https://doi.org/10.1016/j.procbio.2019.05.023>
- [104] Kale C, Gökçek M (2020) A techno-economic assessment of landfill gas emissions and energy recovery potential of different landfill areas in Turkey. *J Clean Prod* 275:122946. <https://doi.org/10.1016/j.jclepro.2020.122946>
- [105] Kargbo H et al (2021) “Drop-in” fuel production from biomass: critical review on techno-economic feasibility and sustainability. *Renew Sustain Energy Rev* 135:110168. <https://doi.org/10.1016/j.rser.2020.110168>
- [106] Karić N et al (2022) Bio-waste valorisation: agricultural wastes as biosorbents for removal of (in)organic pollutants in wastewater treatment. *Chem Eng J Adv* 9:100239. <https://doi.org/10.1016/j.cej.2021.100239>
- [107] Kaya M (2016) Recovery of metals and nonmetals from electronic waste by physical and chemical recycling processes. *Waste Manage* 57:64–90. <https://doi.org/10.1016/j.wasman.2016.08.004>

- [108]Kaza S et al (2018) What a waste 2.0: a global snapshot of solid waste management to 2050. World Bank Publications. <https://datatopics.worldbank.org/what-a-waste/>
- [109]Keng ZX et al (2020) Community-scale composting for food waste: a life-cycle assessment-supported case study. J Clean Prod 261:121220. <https://doi.org/10.1016/j.jclepro.2020.121220>
- [110]Khan MM-U-H et al (2018) Optimal siting of solid waste-to-value-added facilities through a GIS-based assessment. Sci Total Environ 610:1065–1075. <https://doi.org/10.1016/j.scitotenv.2017.08.169>
- [111]Khan S et al (2022a) Technologies for municipal solid waste management: current status, challenges, and future perspectives. Chemosphere 288:132403. <https://doi.org/10.1016/j.chemosphere.2021.132403>
- [112]Khan A et al (2022b) Bio-organics management: novel strategies to manage root-knot nematode, *meloidogyne incognita* pest of vegetable crops. Gesunde Pflanzen. <https://doi.org/10.1007/s10343-022-00679-2>
- [113]Khanal SK et al (2021) Anaerobic digestion beyond biogas. Bioresource Technol 337:125378. <https://doi.org/10.1016/j.biortech.2021.125378>
- [114]Khosravi A et al (2022) Production and characterization of hydrochars and their application in soil improvement and environmental remediation. Chem Eng J 430:133142. <https://doi.org/10.1016/j.cej.2021.133142>
- [115]Kizito S et al (2019) Role of nutrient-enriched biochar as a soil amendment during maize growth: exploring practical alternatives to recycle agricultural residuals and to reduce chemical fertilizer demand. Sustainability 11:3211. <https://doi.org/10.3390/su11113211>
- [116]Koskinopoulou M et al (2021) Robotic waste sorting technology: toward a vision-based categorization system for the industrial robotic separation of recyclable waste. IEEE Robot Autom Mag 28:50–60. <https://doi.org/10.1109/MRA.2021.3066040>
- [117]Kostas ET et al (2017) The application of microwave heating in bioenergy: a review on the microwave pre-treatment and upgrading technologies for biomass. Renew Sustain Energy Rev 77:12–27. <https://doi.org/10.1016/j.rser.2017.03.135>
- [118]Kostas ET et al (2020) Microwave pyrolysis of olive pomace for bio-oil and bio-char production. Chem Eng J 387:123404. <https://doi.org/10.1016/j.cej.2019.123404>
- [119]Kulkarni BN (2020) Environmental sustainability assessment of land disposal of municipal solid waste generated in Indian cities—a review. Environ Dev 33:1–13. <https://doi.org/10.1016/j.envdev.2019.100490>
- [120]Kumar N, Gupta SK (2021) Exploring the feasibility of thermal digestion process: a novel technique, for the rapid treatment and reuse of solid organic waste as organic fertilizer. J Clean Prod 318:128600. <https://doi.org/10.1016/j.jclepro.2021.128600>
- [121]Kumla J et al (2020) Cultivation of mushrooms and their lignocellulolytic enzyme production through the utilization of agro-industrial waste. Molecules 25:2811. <https://doi.org/10.3390/molecules25122811>
- [122]Lee U et al (2017) Evaluation of landfill gas emissions from municipal solid waste landfills for the life-cycle analysis of waste-to-energy pathways. J Clean Prod 166:335–342. <https://doi.org/10.1016/j.jclepro.2017.08.016>
- [123]Leno N et al (2021) Thermochemical digestate fertilizer from solid waste: characterization, labile carbon dynamics, dehydrogenase activity, water holding capacity and biomass allocation in banana. Waste Manag 123:1–14. <https://doi.org/10.1016/j.wasman.2021.01.002>
- [124]Li J et al (2021a) Influence of industrial solid waste as filling material on mechanical and microstructural characteristics of cementitious backfills. Constr Build Mater 299:124288. <https://doi.org/10.1016/j.conbuildmat.2021.124288>
- [125]Li J et al (2021b) Land space simulation of urban agglomerations from the perspective of the symbiosis of urban development and ecological protection: a case study of Changsha-Zhuzhou-Xiangtan urban agglomeration. Ecol Ind 126:107669. <https://doi.org/10.1016/j.ecolind.2021.107669>
- [126]Liu Z et al (2013) Production of solid biochar fuel from waste biomass by hydrothermal carbonization. Fuel 103:943–949. <https://doi.org/10.1016/j.fuel.2012.07.069>
- [127]Liu L et al (2021) Excessive application of chemical fertilizer and organophosphorus pesticides induced total phosphorus loss from planting causing surface water eutrophication. Sci Rep 11:1–8. <https://doi.org/10.1038/s41598-021-02521-7>
- [128]Longsheng C et al (2022) An integrated SWOT-multi-criteria analysis of implementing sustainable waste-to-energy in Pakistan. Renew Energy 195:1438–1453. <https://doi.org/10.1016/j.renene.2022.06.112>
- [129]Lu X, Guo Y (2019) Removal of Pb (II) from aqueous solution by sulfur-functionalized walnut shell. Environ Sci Pollut Res 26:12776–12787. <https://doi.org/10.1007/s11356-019-04753-7>
- [130]Lu K et al (2021) Integration of life cycle assessment and life cycle cost using building information modeling: a critical review. J Clean Prod 285:125438. <https://doi.org/10.1016/j.jclepro.2020.125438>
- [131]Luo M et al (2018) A novel modification of lignin on corn-cob-based biochar to enhance removal of cadmium from water. Biores Technol 259:312–318. <https://doi.org/10.1016/j.biortech.2018.03.075>
- [132]Mabalane PN et al (2021) A techno-economic analysis of anaerobic digestion and gasification hybrid system: energy recovery from municipal solid waste in South Africa. Waste Biomass Valorization 12:1167–1184. <https://doi.org/10.1007/s12649-020-01043-z>
- [133]Mahmud R et al (2021) Integration of techno-economic analysis and life cycle assessment for sustainable process design—a review. J Clean Prod 317:128247. <https://doi.org/10.1016/j.jclepro.2021.128247>
- [134]Maleita C et al (2017) Naphthoquinones from walnut husk residues show strong nematocidal activities against the root-knot nematode *meloidogyne hispanica*. ACS Sustain Chem Eng 5:3390–3398. <https://doi.org/10.1021/acssuschemeng.7b00039>
- [135]Mandal K (2019) Review on evolution of municipal solid waste management in India: practices, challenges and policy implications. J Mater Cycles Waste Manage 21:1263–1279. <https://doi.org/10.1007/s10163-019-00880-y>
- [136]Maroušek J et al (2020) Techno-economic assessment of potato waste management in developing economies. Clean Technol Environ Policy 22:937–944. <https://doi.org/10.1007/s10098-020-01835-w>
- [137]Mavridis S, Voudrias EA (2021) Using biogas from municipal solid waste for energy production: comparison between anaerobic digestion and sanitary landfilling. Energy Convers Manage 247:114613. <https://doi.org/10.1016/j.enconman.2021.114613>
- [138]Md Badrul Hisham NH et al (2019) Production of biosurfactant produced from used cooking oil by *Bacillus* sp. HIP3 for heavy metals removal. Molecules 24:2617. <https://doi.org/10.3390/molecules24142617>
- [139]Meneguzzo F et al (2019) Real-scale integral valorization of waste orange peel via hydrodynamic cavitation. Processes 7:581. <https://doi.org/10.3390/pr7090581>
- [140]Mian MM et al (2017) Municipal solid waste management in China: a comparative analysis. J Mater Cycles Waste Manage 19:1127–1135. <https://doi.org/10.1007/s10163-016-0509-9>

- [141]Mishra S et al (2021) The utilization of agro-biomass/byproducts for effective bio-removal of dyes from dyeing wastewater: a comprehensive review. *J Environ Chem Eng* 9:104901. <https://doi.org/10.1016/j.jece.2020.104901>
- [142]Mlaik N et al (2019) Enzymatic pre-hydrolysis of organic fraction of municipal solid waste to enhance anaerobic digestion. *Biomass Bioenerg* 127:105286. <https://doi.org/10.1016/j.biombioe.2019.105286>
- [143]Moharm AE et al (2022) Fabrication and characterization of effective biochar biosorbent derived from agricultural waste to remove cationic dyes from wastewater. *Polymers* 14:2587. <https://doi.org/10.3390/polym14132587>
- [144]Monda H et al (2017) Molecular characteristics of water-extractable organic matter from different composted biomasses and their effects on seed germination and early growth of maize. *Sci Total Environ* 590:40–49. <https://doi.org/10.1016/j.scitotenv.2017.03.026>
- [145]Mondal P et al (2021) Municipal solid waste fired combined cycle plant: techno-economic performance optimization using response surface methodology. *Energy Convers Manage* 237:114133. <https://doi.org/10.1016/j.enconman.2021.114133>
- [146]Moult J et al (2018) Greenhouse gas emissions of food waste disposal options for UK retailers. *Food Policy* 77:50–58. <https://doi.org/10.1016/j.foodpol.2018.04.003>
- [147]Mourad M (2016) Recycling, recovering and preventing “foodwaste”: competing solutions for food systems sustainability in the United States and France. *J Clean Pro* 126:461–477. <https://doi.org/10.1016/j.jclepro.2016.03.084>
- [148]Mr P et al (2022) Recycling of agricultural (orange and olive) biowastes into ecofriendly fertilizers for improving soil and garlic quality. *Resour, Conserv Recycling Adv* 15:200083. <https://doi.org/10.1016/j.rcradv.2022.200083>
- [149]Mrozik W et al (2021) Environmental impacts, pollution sources and pathways of spent lithium-ion batteries. *Energy Environ Sci* 14:6099–6121. <https://doi.org/10.1039/D1EE00691F>
- [150]Murtaza B et al (2019) Municipal solid waste compost improves crop productivity in saline-sodic soil: a multivariate analysis of soil chemical properties and yield response. *Commun Soil Sci Pla* 50:1013–1029. <https://doi.org/10.1080/00103624.2019.1603305>
- [151]Myers SS et al (2017) [Accepted Manuscript] climate change and global food systems: potential impacts on food security and undernutrition. *Annu Rev Public Health*. <https://doi.org/10.1146/annurev-publhealth-031816-044356>
- [152]Nabavi V et al (2020) Feasibility study on the production and consumption of wood pellets in Iran to meet return-on investment and greenhouse gas emissions targets. *Renew Energy* 151:1–20. <https://doi.org/10.1016/j.renene.2019.10.140>
- [153]Ng KS et al (2021) Techno-economic assessment of a novel integrated system of mechanical-biological treatment and valorisation of residual municipal solid waste into hydrogen: a case study in the UK. *J Clean Prod* 298:126706. <https://doi.org/10.1016/j.jclepro.2021.126706>
- [154]Norouzi O, Dutta AJE (2022) The current status and future potential of biogas production from Canada’s organic fraction municipal solid waste. *Energies* 15:475. <https://doi.org/10.3390/en15020475>
- [155]Obeng GY et al (2020) Coconut wastes as bioresource for sustainable energy: quantifying wastes, calorific values and emissions in Ghana. *Energies* 13:2178. <https://doi.org/10.3390/en13092178>
- [156]Ogunjuyigbe A et al (2017) Electricity generation from municipal solid waste in some selected cities of Nigeria: an assessment of feasibility, potential and technologies. *Renew Sustain Energy Rev* 80:149–162. <https://doi.org/10.1016/j.rser.2017.05.177>
- [157]Olujobi O et al (2022) Conversion of organic wastes to electricity in Nigeria: legal perspective on the challenges and prospects. *Int J Environ Sci Te* 19:939–950. <https://doi.org/10.1007/s13762-020-03059-3>
- [158]Oni BA et al (2019) Significance of biochar application to the environment and economy. *Annals Agric Sci* 64:222–236. <https://doi.org/10.1016/j.aos.2019.12.006>
- [159]Osman AI et al (2022a) Hydrogen production, storage, utilisation and environmental impacts: a review. *Environ Chem Lett* 20:153–188. <https://doi.org/10.1007/s10311-021-01322-8>
- [160]Osman AI et al (2022b) Biochar for agronomy, animal farming, anaerobic digestion, composting, water treatment, soil remediation, construction, energy storage, and carbon sequestration: a review. *Environ Chem Lett* 20:2385–2485. <https://doi.org/10.1007/s10311-022-01424-x>
- [161]Otsuka K, Fan S (2021) Agricultural development: new perspectives in a changing world. International Food Policy Research Institute, Washington, D.C.
- [162]Pandey A, Asif M (2022) Assessment of energy and environmental sustainability in South Asia in the perspective of the sustainable development goals. *Renew Sustain Energy Rev* 165:112492. <https://doi.org/10.1016/j.rser.2022.112492>
- [163]Pardo-Giménez A et al (2020) Optimization of cultivation techniques improves the agronomic behavior of *Agaricus subrufescens*. *Sci Rep* 10:1–9. <https://doi.org/10.1038/s41598-020-65081-2>
- [164]Patel SK et al (2021) Integrating strategies for sustainable conversion of waste biomass into dark-fermentative hydrogen and value-added products. *Renew Sustain Energy Rev* 150:111491. <https://doi.org/10.1016/j.rser.2021.111491>
- [165]Patrizio P et al (2015) Biomethane as transport fuel—a comparison with other biogas utilization pathways in northern Italy. *Appl Energy* 157:25–34. <https://doi.org/10.1016/j.apenergy.2015.07.074>
- [166]Pawel I (2014) The cost of storage—how to calculate the levelized cost of stored energy (LCOE) and applications to renewable energy generation. *Energy Procedia* 46:68–77. <https://doi.org/10.1016/j.egypro.2014.01.159>
- [167]Peng C et al (2016) Production of char from sewage sludge employing hydrothermal carbonization: char properties, combustion behavior and thermal characteristics. *Fuel* 176:110–118. <https://doi.org/10.1016/j.fuel.2016.02.068>
- [168]Petrillo A et al (2022) Multi-criteria analysis for life cycle assessment and life cycle costing of lightweight artificial aggregates from industrial waste by double-step cold bonding palletization. *J Clean Prod* 351:131395. <https://doi.org/10.1016/j.jclepro.2022.131395>
- [170]Picot-Allain C et al (2021) Conventional versus green extraction techniques—a comparative perspective. *Curr Opin Food Sci* 40:144–156. <https://doi.org/10.1016/j.cofs.2021.02.009>
- [171]Porter C (2016). World agricultural prospects the road to 2050. Supply Intelligence Ltd.
- Praveen S et al (2021) Techno-economic feasibility of biochar as biosorbent for basic dye sequestration. *J Indian Chem Soc* 98:100107. <https://doi.org/10.1016/j.jics.2021.100107>
- [172]Pryshlakivsky J, Searcy C (2021) Life cycle assessment as a decisionmaking tool: practitioner and managerial considerations. *J Clean Prod* 309:127344. <https://doi.org/10.1016/j.jclepro.2021.127344>

- [173]Pu X et al (2020) Utilization of industrial waste lithium-silicon-powder for the fabrication of novel nap zeolite for aqueous Cu (II) removal. *J Clean Prod* 265:121822. <https://doi.org/10.1016/j.jclepro.2020.121822>
- [174]Rajeh C et al (2021) Food loss and food waste recovery as animal feed:a systematic review. *J Matr Cycles Wast* 23:1–17. <https://doi.org/10.1007/s10163-020-01102-6>
- [175]Rashid MI, Shahzad K (2021) Food waste recycling for compost production and its economic and environmental assessment as circular economy indicators of solid waste management. *J Clean Pro* 317:128467. <https://doi.org/10.1016/j.jclepro.2021.128467>
- [176]Razzaq A et al (2021) Dynamic and causality interrelationships from municipal solid waste recycling to economic growth, carbon emissions and energy efficiency using a novel bootstrapping autoregressive distributed lag. *Resource Conserve Recycle* 166:105372. <https://doi.org/10.1016/j.resconrec.2020.105372>
- [177]Rittl TF et al (2018) Greenhouse gas emissions from soil amended with agricultural residue biochars: effects of feedstock type, production temperature and soil moisture. *Biomass Bioenergy* 117:1–9. <https://doi.org/10.1016/j.biombioe.2018.07.004>
- [178]Roman FF et al (2021) Hydrochars from compost derived from municipal solid waste: production process optimization and catalytic applications. *J Environ Chem Eng* 9:104888. <https://doi.org/10.1016/j.jece.2020.104888>
- [179]Rosa-Clot M, Tina GM (2020) Chapter 10–levelized cost of energy(LCOE) analysis. In: Rosa-Clot M, Marco Tina G (eds) *Floating PV plants*. Academic Press, pp 119–127. <https://doi.org/10.1016/B978-0-12-817061-8.00010-5>
- [180]Rosales E et al (2017) Challenges and recent advances in biochar as low-cost biosorbent: from batch assays to continuous-flow systems. *Biorsource Technol* 246:176–192. <https://doi.org/10.1016/j.biortech.2017.06.084>
- [181]Rybacka J et al (2016) Technology readiness level assessment of composites recycling technologies. *J Clean Prod* 112:1001–1012. <https://doi.org/10.1016/j.jclepro.2015.08.104>
- [182]Saad S et al (2010) Chemically modified sugarcane bagasse as a potentially low-cost bio sorbent for dye removal. *Desalination* 264:123–128. <https://doi.org/10.1016/j.desal.2010.07.015>
- [183]Sadeghi S et al (2022) Microbial characteristics of municipal solid waste compost: occupational and public health risks from surface applied compost. *Waste Manage* 144:98–105. <https://doi.org/10.1016/j.wasman.2022.03.012>
- [184]Saja AMA et al (2021) Municipal solid waste management practices and challenges in the south eastern coastal cities of Sri Lanka. *Sustainability* 13:4556. <https://doi.org/10.3390/su13084556>
- [185]Salemdeeb R et al (2017) Environmental and health impacts of using food waste as animal feed: a comparative analysis of food waste management options. *J Clean Prod* 140:871–880. <https://doi.org/10.1016/j.jclepro.2016.05.049>
- [186]Solis M, Silveira S (2020) Technologies for chemical recycling of household plastics—a technical review and TRL assessment. *Waste Manage* 105:128–138. <https://doi.org/10.1016/j.wasman.2020.01.038>
- [187]Sun C et al (2021) Compound utilization of construction and industrial waste as cementitious recycled powder in mortar. *Resour Conserve Recycle* 170:105561. <https://doi.org/10.1016/j.resconrec.2021.105561>
- [188]Tahat M et al (2020) Soil health and sustainable agriculture. Excessive and disproportionate use of chemicals cause soil contamination and nutritional stress 12:4859. <https://doi.org/10.3390/su12124859>
- [189]Talwar N, Holden NM (2022) The limitations of bioeconomy LCA studies for understanding the transition to sustainable bioeconomy. *Int J Life Cycle Assess* 27:680–703. <https://doi.org/10.1007/s11367-022-02053-w>
- [190]Tawfk A et al (2022) Methods to alleviate the inhibition of sludge anaerobic digestion by emerging contaminants: a review. *Environ Chem Lett*. <https://doi.org/10.1007/s10311-022-01465-2>
- [191]Thanigaivel S et al (2022) Exploration of effective biorefinery approach to obtain the commercial value-added products from algae. *Sustain Energy Technol Assess* 53:102450. <https://doi.org/10.1016/j.seta.2022.102450>
- [192]Timilsina GR (2021) Are renewable energy technologies cost competitive for electricity generation? *Renew Energy* 180: 658–672. <https://doi.org/10.1016/j.renene.2021.08.088>
- [193]Torok VA et al (2021) Human food waste to animal feed: opportunities and challenges. *Anim Prod Sci* 2:1129–1139. <https://doi.org/10.1071/AN20631>
- [194]Tripathi N et al (2019) Biomass waste utilisation in low-carbon products: harnessing a major potential resource. *NPJ Clim Atmos Sci* 2:1–10. <https://doi.org/10.1038/s41612-019-0093-5>
- [195]Tun MM, Juchelkova D (2018) Assessment of solid waste generation and greenhouse gas emission potential in Yangon city, Myanmar. *J Mater Cycles Waste Manage* 20:1397–1408. <https://doi.org/10.1007/s10163-017-0697-y>
- [196]Tyagi VK et al (2018) Anaerobic co-digestion of organic fraction of municipal solid waste (OFMSW): progress and challenges. *Renew Sustain Energy Rev* 93:380–399. <https://doi.org/10.1016/j.rser.2018.05.051>
- [197]Uddin M et al (2021) Prospects of bioenergy production from organic waste using anaerobic digestion technology: a mini review. *Front Energy Res* 9:627093. <https://doi.org/10.3389/ferg.2021.627093>
- [198]Usmani Z et al (2020) Advancement in valorization technologies to improve utilization of bio-based waste in bioeconomy context. *Renew Sust Energy Rev* 131:109965. <https://doi.org/10.1016/j.rser.2020.109965>
- [199]Van Nguyen TT et al (2022) Valorization of agriculture waste biomass as biochar: as first-rate biosorbent for remediation of contaminated soil. *Chemosphere* 307:135834. <https://doi.org/10.1016/j.chemosphere.2022.135834>
- [200]Wang C et al (2022) Reduction in net greenhouse gas emissions through a combination of pig manure and reduced inorganic fertilizer application in a double-rice cropping system: three-year results. *Agric Ecosyst Environ* 326:107799. <https://doi.org/10.1016/j.agee.2021.107799>
- [201]Wiśniewska P et al (2022) Waste tire rubber devulcanization technologies: state-of-the-art, limitations and future perspectives. *Waste Manage* 150:174–184. <https://doi.org/10.1016/j.wasman.2022.07.002>
- [202]Wu F et al (2022) High value-added resource utilization of solid waste: review of prospects for supercritical CO<sub>2</sub> extraction of valuable metals. *J Clean Prod* 372:133813. <https://doi.org/10.1016/j.jclepro.2022.133813>
- [203]Xiao L et al (2021) Biochar promotes methane production during anaerobic digestion of organic waste. *Environ Chem Lett* 19:3557–3564. <https://doi.org/10.1007/s10311-021-01251-6>
- [204]Xiu S, Shahbazi A (2012) Bio-oil production and upgrading research:a review. *Renew Sustain Energy Rev* 16:4406–4414. <https://doi.org/10.1016/j.rser.2012.04.028>

- [205]Xue X et al (2022) Thermodynamic and economic analyses of a new compressed air energy storage system incorporated with a waste-to-energy plant and a biogas power plant. *Energy* 261:125367. <https://doi.org/10.1016/j.energy.2022.125367>
- [206]Yaashikaa PR, Kumar PS (2022) Bioremediation of hazardous pollutants from agricultural soils: a sustainable approach for waste management towards urban sustainability. *Environ Pollut* 312:120031. <https://doi.org/10.1016/j.envpol.2022.120031>
- [207]Yan Y et al (2022) Enhancing enzyme activity via low-intensity ultrasound for protein extraction from excess sludge. *Chemosphere* 303:134936. <https://doi.org/10.1016/j.chemosphere.2022.134936>
- [208]Yang N et al (2012) Greenhouse gas emissions from MSW incineration in China: impacts of waste characteristics and energy recovery. *Waste Manage* 32:2552–2560. <https://doi.org/10.1016/j.wasman.2012.06.008>
- [209]Yang Y et al (2019) Estimate of restaurant food waste and its biogas production potential in China. *J Clean Pro* 211:309–320. <https://doi.org/10.1016/j.jclepro.2018.11.160>
- [210]Yang M et al (2022) Circular economy strategies for combating climate change and other environmental issues. *Environ Chem Lett*. <https://doi.org/10.1007/s10311-022-01499-6>
- [211]Yelatontsev D (2023) Production of versatile biosorbent via ecofriendly utilization of non-wood biomass. *Chem Eng J* 451:138811. <https://doi.org/10.1016/j.cej.2022.138811>
- [212]Yong ZJ et al (2021) Biogas and biofertilizer production from organic fraction municipal solid waste for sustainable circular economy and environmental protection in Malaysia. *Sci Total Environ* 776:145961. <https://doi.org/10.1016/j.scitotenv.2021.145961>
- [213]Zhao J et al (2022) How renewable energy alleviate energy poverty? A global analysis. *Renew Energy* 186:299–311. <https://doi.org/10.1016/j.renene.2022.01.005>
- [214]Zhou Z, Zhang L (2022) Sustainable waste management and waste to energy: valuation of energy potential of MSW in the Greater Bay Area of China. *Energy Policy* 163:112857. <https://doi.org/10.1016/j.enpol.2022.112857>
- [215]Zhou H et al (2017) Water sources of *Nitraria sibirica* and response to precipitation in two desert habitats. *J Appl Ecol* 28:2083–2092. <https://doi.org/10.13287/j.1001-9332.201707.021>
- [216]Zhu Y et al (2021) A review of municipal solid waste in China: characteristics, compositions, influential factors and treatment technologies. *Environ Dev Sustain* 23:6603–6622. <https://doi.org/10.1007/s10668-020-00959-9>
- [217]Zu Ermgassen EK et al (2016) Reducing the land use of EU pork production: where there's swill, there's a way. *Food Policy* 58:35–48. <https://doi.org/10.1016/j.foodpol.2015.11.001>

