

# Pharmacognostic Study, Extraction, Formulation And Evaluation Of Polyherbal Gel

Pradeep P. Sonwane<sup>1\*</sup>, Vaishnavi Nikam<sup>2</sup>, Hitisha Kadu<sup>3</sup>, Chaitnaya Durge<sup>4</sup>, Rushubh Kothari<sup>5</sup>

<sup>1</sup>Assitant Professor, <sup>2</sup>Student, <sup>3</sup>Student, <sup>4</sup>Student, <sup>5</sup>Student

AISSMS College Of Pharmacy, Pune 411001 Maharashtra, India

**Abstract** - Substances derived from plants and herbal therapy have attracted the attention of customers due to the presence of bioactive compounds and the least side-effects. *Tridax procumbens* have been used as an excellent anti-inflammatory and wound-healing drug. The other active pharmaceutical component of this formulation is *Murraya Koenigii* which has the ability to show anti-inflammatory action. Keeping these things in mind, we have formulated poly-herbal gel. Extraction and evaluation of the crude drug were carried out. The hydro-alcoholic extract was used for the formulation. After the completion of the polyherbal formulation, it was evaluated with its physicochemical studies like color, pH, Spreadability, consistency, etc. The stability of the formulation was evaluated at various temperature conditions which shows no change in the irritancy, or spreadability.

**Keywords:** *Tridax Procumbens*, *Murraya Koenigii*, Topical Gel, Polyherbal Gel, Brookfield viscometer, anti-inflammatory

## INTRODUCTION:

Polyherbal Gel the name itself explains that it is made of more than two drugs this polyherbal formulation is made from two herbs which is *Tridax procumbens* and *Murraya Koenigii* which is known for its antioxidant and anti-inflammatory action. <sup>(1)</sup>

Traditional Drugs and indispensable drugs are both heavily reliant on herbal remedies, both in developed and developing nations. The supposition that plants are safe, secure, and have minimal side effects has conducted considerable interest in medications produced from them. A review of the literature reveals that traditional plant medications are helpful for several skin-related problems and for crack mending. World Health Organization (WHO) as well as our country has been elevating the use of traditional drugs because they are less precious, easily available, and comprehensive, especially in developing countries.

The best thing about using herbal drug is that they can be applied by people of any age group and has no side goods or ineffective cures. Multiple explorations have used the extracts of *Tridax procumbens* L., which belongs to the Asteraceae family, and *Murraya Koenigii* (curry leaves), which belongs to the Rutaceae family.

For a variety of skin conditions, further than 80 of the world's population still uses traditional treatments. Herbal remedies for crack care involve cleaning, debridement, and creating a wettish terrain to promote the creation of the ideal conditions for the natural mending process. Literature reveals that simple traditional shops (*Adhatoda vasica*, *Boerhaavia diffusa*, *Caesalpinia sappan*, *Curcuma longa*, *Cyperus rotundus*, *Eclipta alba*, *Glycyrrhiza glabra*, *Gymnema sylvestre*, *Oryza sativa*, *Piper longum*, *Plumbago zeylanica*) are helpful in the treatment of several skin-related problems.

The present exploration reflected exploring the presumptive part(s) of active excerpts of *Tridax procumbens* (whole factory), *Murraya Koenigii* (leaves), *Calendula officinalis*(flowers), and *Aloe barbadensis*(leaves) formulated as ointment product. The formulations were characterized by determining the pharmaceutical extrudability, swelling indicator, and washability <sup>(1-2)</sup>

## II. LITERATURE SURVEY

### 1] *Tridax procumbens*:

**ACTIVITY FOUND:** <sup>(3)</sup>

1. Antiviral
2. Anti-oxidant
3. Antibiotic
4. Insecticidal
5. Anti-inflammatory
6. Hair Tonic
7. Anti-Diabetic
8. Anti-Fungal

- **Kingdom**-Plantae
- **Family**- Asteraceae
- **Genus**- Tridax
- **Species**-T. Procumbens

**Common Name**- A) Coatbuttons

B) Tridex Daisy

**Chemical constituents-**

The plant exhibits a variety of chemical components, including methyl, 14-oxoacagacunoate, 14-oxonacosanoate, 3-methyl-non dodecylbenzene, heptacosanyl cyclohexane carboxylate, 12-hydroxytetracos-15-one, -amyrone, fecosterol, and sitosterol, as well as arachidic.

2] **Murraya Koenigii (Linn.) Spreng.**

**ACTIVITY FOUND:** <sup>(4)</sup>

1. Anti-Pyretic
2. Hypoglycaemic
3. Anti-Microbial
4. Hepatoprotection
5. Anti-inflammatory
6. Cytotoxic

**Kingdom**-Plantae

**Family**-Rutaceae

**Genus**-Murraya

**Species**-M. Koenigii

**Common Name**-A) Sweet Neem

B) Curry Leaf Tree

C) Karwa Pale

D) Karapincha

**Chemical constituents-**

Leaves contain essential oil [ 0.14 to 0.32 percent (v/w)] whose main constituents are mono and sesquiterpenes. The major constituents of the essential oil are sabinene (31.8 to 44.8 percent )  $\beta$ - caryophyllene ( 24.4 percent ) ,  $\alpha$ - pinene ( 19 to 19.7 percent ) and  $\beta$ - Phellandrene.

**III. MATERIAL METHOD**

**A) Collection of Plant**

**1) Tridax procumbens- (FIG – 1)**

- Tridax procumbens L. leaves were gathered from various locations in Pune and its environs and properly cleaned with distilled water.
- After the cleaned plant pieces have dried completely in the shade, they are ground into a fine powder using a mechanical grinder and placed in an airtight container.

**2) Murraya Koenigii- (FIG – 2)**

- The plant Murraya Koenigii was collected from the agricultural areas of Pune.
- The fresh leaves were collected and separated from the plant and used for extraction <sup>(6)</sup>

FIG. 1



FIG. 2



#### B) AUTHENTICATION

The collected Plant of *Murraya Koenigii* and *Tridax procumbens* were collected from Pune district and authenticated by Botanical Survey of India, Pune.

#### C) PREPARATION OF EXTRACT

##### 1. *Tridax Procumbens* <sup>(5)</sup>

- The collected material that is leaves of *Tridax Procumbens* were shade dried and powdered.
- The powder was macerated in ethanol and water which was taken in ratio of 30:70 for 24 hours.
- The solution is filtered with Whatman filter paper. This solution was the evaporated-on room temperature to get the desired extract and TLC was performed to analyse the constituents.
- The extract was then further used for making the formulation.

##### 2. *Murraya Koenigii* <sup>(6)</sup>

- The powder of leaves *Murraya Koenigii* was macerated in water and methanol.
- The prepared solution was then filtered after 24 hours using Whatman filter paper.
- This solution was the evaporated-on room temperature to get the desired extract and TLC was performed for investigating the constituents present in the extract. This extract was then further used for making the formulation.

#### D) PREPARATION OF GEL

##### Instrument:

- The instrument for formulating or mixing the gel was Mechanical Stirrer of Remi Motor Type nQ-122.
- The speed for stirrer was set of 500-600 RPM for proper mixing.

##### Preparation of Carbopol: (FIG – 9)

- Carbopol was soaked in water overnight for making the gel.
- The quantity of water was 1/4<sup>th</sup> of the total quantity which was to be mixed in the whole formulation.
- As after the soaking the Carbopol swells which makes it easier for uniform mixing.



Color and appearance of Carbopol 934: (Table 1)  
Table - 1

TEST	RESULT
Color	White
Appearance	Powder



FIG – 9

**Procedure for Gel:**

1. The soaked Carbopol which was prepared were mixed with the help of mechanical stirrer for making the homogenised mixture.
2. Additional water was added for mixing the gel and the drug extracts of Tridax Procumbens and Murraya Koenigii in the ratio of 1:1.
3. Addition of Triethylamine was done for getting the desired viscosity and pH of the Gel.
4. Triethanolamine was added for adjusting the pH of gel for with skin pH
5. In last preservatives methyl paraben and propyl paraben was added. <sup>(8,9)</sup>
- 6.

Sr. no.	Ingredient	Quantity
1.	Carbopol - 934	2 g
2.	Tridax Procumbens	0.5 g
3.	Murraya Koenigii	0.5 g
4.	Triethylamine	0.1 ml
5.	Methyl Paraben	0.1 g
6.	Propyl Paraben	0.05 g
7.	Water	q.s to 100ml

### III. EVALUATION

#### A) Crude Drug

##### 1. TLC of the Powder:

##### Murraya Koenigii

1. With the help of TLC preliminary test was performed for the conformation of constituents in the extract.
2. Mobile phase as Toluene and ethyl acetate in the ratio of 3:7 for the separation was taken.
3. Three mixtures were taken for mixing the drug Benzene, Water, and alcohol i.e., methanol and best separation and presence of alkaloid was found in methanol and hence preparation of extracts was done in alcohol. <sup>(11)</sup>

##### B) Tridax Procumbens

1. Thin layer chromatography for Tridax Procumbens for Beta-sitosterol developed at 25 + 2 degree.
2. Mobile Phase Chloroform, Methanol, acetic acid in ratio 8:0.5:0.5 V/V as mobile phase in an image.
3. Then developed 8 cm then dried.

Derivatized with anisaldehyde is spraying agent (0.2ml Anisaldehyde +20 ml acetic acid + Concentrated 0.5 ml Sulphuric acid) in the derivatising chamber for 20 min again dried it. After drying heated 105 degrees Celsius for 10 min.

##### 3. Ash Value <sup>(11-12)</sup>

The residue remaining after incineration is the ash content of the medicine. Contains inorganic salts of carbonates, phosphates, and silicates of Na, K, Ca, and Mg. A high ash value indicates impurity, contamination, or neglectfulness in preparing crude medicine.

Methods to determine ash value:

##### Total Ash

- Measures the total amount of material remaining after ignition.
- Physiological ash: plant-derived
- Non-physiological ash: sand/soil adhered to plant tissue.
- %Total ash=100\* Wash/Wdr

##### Water soluble ash

- Difference in weight between total ash & residue after treatment of total ash with water.

##### Insoluble ash

- Residue obtained after boiling total ash + dil. HCl and igniting remaining insoluble matter. Measure the amount of silica present.

#### B) Formulation:

- 1) **Physical Evaluation:** Aspects of the body, such as colour and look, were assessed.
- 2) **Homogeneity:** After being put in the container, all generated gels were visually inspected for uniformity for existence of any aggregate.
- 3) **pH:** A digital pH meter was used to determine the pH of different gel compositions. A precise 2.5gm of gel was weighed, mixed with 25ml of purified water, and kept for two hours. Three samples of each formulation's pH were used to measure it, and the average results are shown. Using a pH meter, the pH of the dispersions was determined.
- 4) **Viscosity:** Herbal gel's viscosity was measured using a Brookfield rotating viscometer at 50 rpm with spindle no. 64. At the conclusion of the two minutes, when the sample had reached equilibrium, each reading was obtained. Three samples' viscosities were determined three times.  
Herbal gel's viscosity was measured using a Brookfield rotating viscometer at 50 rpm with spindle no. 64. At the conclusion of the two minutes, when the sample had reached equilibrium, reading was obtained.
- 5) **Irritability:** Human subjects' skin was exposed to test formulations, and the results were tracked.
- 6) **Stability analysis:** The formulations underwent a four-week physical stability test at a range of temperatures, including 2°C and 25°C. Within four weeks, it was discovered that the formulation was physically stable at various temperatures, such as 2°C and 25°C

IV. RESULTS

A) Crude Drug

1. TLC

	Murraya Koenigii	Tridax Procumbens
<b>Rf Values</b>	<b>0.8</b>	<b>0.5</b>

2. Ash Value <sup>(11-12)</sup>

Sr. No	Particular	Murraya Koenigii	Standard	Tridax Procumbens	Standard
1.	Total Ash	7.2%	7.2%	18%	20.4%
2.	Water Soluble Ash	9%	10.4%	6%	6.5%
3.	Acid Insoluble Ash	18%	35.6%	6%	7.1%

3. Preliminary Phyto-Chemical Screening

Sr.no	Test	Tridax Procumbens	Murraya Koenigii
1)	Test for carbohydrates	(+ve)	(+ve)
2)	Test for steroids	(+ve)	(+ve)
3)	Test for alkaloids	(+ve)	(+ve)
4)	Test for saponins	(+ve)	(+ve)
5)	Test for Flavonoids	(+ve)	(+ve)
6)	Test for Tannin and phenolic compounds	(+ve)	(+ve)

**B) Formulation**

**1. Physical Evaluation**

Sr. no	Particulars	Evaluation
1.	Colour	Greenish-Brown
2.	Appearance	Uniform
3.	Odour	Pleasant
4.	Texture	Smooth

**2. Homogeneity**

Sr.no	Batch	Homogeneity
1.	F1	Homogenous
2.	F2	Homogenous
3.	F3	Homogenous

**3. pH**

Sr no	Batch	pH
1)	F1	6.08
2)	F2	6.1
3)	F3	6.42

**4. Viscosity**

Rpm	Cp
50	36640

**V) DISCUSSION**

Herbal drug has come an item of global significance both medicinal and provident. Although the operation of these herbal drugs has increased, their quality, safety, and effectiveness are serious enterprises in industrialized and developing countries. Herbal remedies are getting adding patient compliance as they're devoid of typical side goods of allopathic drugs. The present exploration has been accepted with the end to formulate and estimate the polyherbal gel containing *Tridax procumbens* splint excerpt and *Murraya Koenigii*. The gel expression was designed by using Carbopol 940, *Tridax procumbens* and *Murraya Koenigii* excerpt, methyl paraben, propyl paraben, and the needed quantum of distilled water. The skin pH was maintained by the dropwise addition of Triethanolamine. The physicochemical parameters of phrasings (pH, spreadibility, density, etc.) were determined. Herbal specifics are considered safer than allopathic drugs as allopathic drugs are associated with side goods. One of the styles for its survival is the medication of excerpts and their phrasings for better immersion and penetration of the active half into the systemic rotation.

**VI) REFERENCE**

1. Prasad, G. B. K. S., Dua, V. K., & Mathur, A. B. H. I. S. H. E. K. (2011). Anti-inflammatory activity of leaves extracts of *Murraya koenigii* L. *Int J Pharm Biol Sci*, 2(1), 541-544.
2. Lokesh Prasad, M. S., Gurunath, K. P., Chandrasekar, S. B., Umashankar, C., & Pawar, A. T. (2017). Formulation and evaluation of herbal formulations (Ointment, Cream, Gel) containing *Tridax procumbens* and *Areca catachu*. *Journal of Scientific and Innovative Research*, 6(3), 97-100.
3. Patel, N. A., Patel, M., & Patel, R. P. (2011). Formulation and evaluation of polyherbal gel for wound healing. *Int Res J Pharm*, 1(1), 15-20.
4. Iyer, D., & Devi, U. (2008). Phyto-pharmacology of *Murraya koenigii* (L.). *Pharmacognosy Reviews*, 2(3), 180.



5. Dattaray, D. (2022). Traditional Uses and Pharmacology of Plant *Tridax procumbens*: A Review. *Sys Rev Pharm*, 13(5), 511-517.
6. Ghosh, P., Biswas, S., Biswas, M., Dutta, A., Sil, S., & Chatterjee, S. (2019). Morphological, Ethno biological and Phytopharmacological Attributes of *Tridax procumbens* Linn. (Asteraceae): A Review. *Int. J. Sci. Res. in Biological Sciences Vol*, 6(2).
7. Nishan, M., & Subramanian, P. (2015). *Murraya koenigii* (curry leave)-A review on its potential. *Int. J. PharmTech Res*, 7(4), 566-572.
8. Varma, M. P. D. K. Therapeutic Potential of *Murraya Koenigii* (Curry Leaves) In Dyslipidemia: A Review.
9. Awasthi, S., Irshad, M., Das, M. K., Ganti, S. S., & Rizvi, M. A. (2009). Anti-inflammatory activity of *Calotropis gigantea* and *Tridax procumbens* on carrageenin-induced paw edema in rats. *Ethnobotanical leaflets*, 2009(5), 2.
10. Rao, L. J. M., Ramalakshmi, K., Borse, B. B., & Raghavan, B. (2007). Antioxidant and radical-scavenging carbazole alkaloids from the oleoresin of curry leaf (*Murraya koenigii* Spreng.). *Food Chemistry*, 100(2), 742-747.
11. Jadhav, V. D., Talele Swati, G., Bakliwal Akshada, A., & Chaudhari, G. N. (2015). Formulation and evaluation of herbal gel containing leaf extract of *Tridax Procumbens*. *J Pharm Biosci*, 3, 65-72.
12. Pande, J., Padalia, H., Donga, S., & Chanda, S. (2018). Pharmacognostic, physicochemical and phytochemical studies of *Andrographis echiodides* Nees. and *Tridax procumbens* L. leaf and stem. *The Pharma Innovation Journal*, 7(6), 303-315.
13. Sandeep, D. S., Nayak, P., Jose, J., Relita, M. R., & Sumana, D. R. (2017). Formulation and evaluation of antibacterial herbal gels of *Murraya koenigii* leaves extract. *Research Journal of Pharmacy and Technology*, 10(6), 1798-1801.
14. Wagh, J. M., Wagh, R., Rathee, K., & Prajapati, C. (2022). Cream Curry Leaf for Dark Spot Reduction. *Saudi J Med Pharm Sci*, 8(5), 234-238.
15. Bansode, P. V., Patil, K. S., & Hajare, A. A. (2020). Bioactivity Guided Antidiabetic Formulation Development of *Tridax procumbens* Linn Leaves. *INDIAN JOURNAL OF PHARMACEUTICAL EDUCATION AND RESEARCH*, 54(3), 705-713.
16. Yadav, S. K., Mishra, M. K., Tiwari, A., & Shukla, A. (2016). Emulgel: a new approach for enhanced topical drug delivery. *Int J Curr Pharm Res*, 9(1), 15-19.
17. Paralkar, S. D., Kamalapurkar, K. A., & Kothari, Y. R. Pharmaceutical Sciences.
18. Sharma, R., & Kumar, U. (2019). Exploration and phytochemical estimation of *Murraya koenigii* leaves for pharmaceutical applications. *Asian Journal of Pharmaceutical Research*, 9(3), 159-168.
19. Asema, S. U. K., Shaikh, A. A., Farooqui, M., & Fatema, S. (2021). Phytochemical analysis of *Murraya koenigii* leaves extract and its biological activity. *The International Journal of Analytical and Experimental Modal Analysis*, 3(1), 1798-1801.
20. Jadhav, V. D., Talele Swati, G., Bakliwal Akshada, A., & Chaudhari, G. N. (2015). Formulation and evaluation of herbal gel containing leaf extract of *Tridax Procumbens*. *J Pharm Biosci*, 3, 65-72.
21. Lohit, R. B., Ramesh, S. S., Ravindra, S. R., Bhausheb, S. S., & Vijay, G. S. FORMULATION AND EVALUATION OF HERBAL GEL USING LEAVES OF TRIDAX PROCUMBENS LINN