

SINGLE BUD PROPAGATION: BOON FOR SUGARCANE INDUSTRY

Raushan Lal

Assistant professor DRMLAU affiliated college, Ayodhya (Faizabad, Uttar Pradesh)

Former SRF, Indian Institute of Sugarcane Research-ICAR, Dilkhusha, Lucknow

Abstract: Sugarcane (*Saccharum Officinarum L.*) is monocotyledon crop plant that mostly propagate through conventional methods (bud chip method, sugarcane initiating technology SSI, Cane node technology). However conventional propagation lacks rapid multiplication. Micropropagation is best alternative of conventional methods. Procedure in Micropropagation is broadly classified as collection of explant, sterilization of explant, inoculation, explant culture, multiplication, rooting, hardening. The variety was cultured on MS medium supplemented with different concentrations of growth regulators on shoot initiation, multiplication, rooting and acclimatisation stages, for initiation stage the best performance was observed on MS medium supplemented 1.0mg/l of BAP. On the other hand multiplication stage was best on MD media enriched with 2.0 mg/l BAP+ 1.0 mg/l NAA, with regard to root induction, the best rooting response achieved best on ½ MS media enriched with 2.0 mg/l NAA +0.5 mg/l BAP. Healthy seeds are a prerequisite for improving the productivity of sugarcane crop. The best way to ensure the production of disease free sugarcane seeds via a technique called 'Meristem Culture' (Tissue culture). According to a study about 80% of seed material can be saved by planting single bud setts over the conventional 3 bud setts planting.

Key words: single bud propagation, Micropropagation, bud chip method, MS medium, Tissue culture.

Introduction: Several technique have been developed for rapid multiplication of seed cane which can be categorised in two groups (a) Biotechnological and

(b) Conventional or traditional method of sett plantation.




Fig 1 : Micropropagation technique

Biotechnological method consist of micropropagation come under tissue culture technique(Salokhe,2021) while conventional methods chiefly include bud chip method,cane node technology (developed by ICAR- IISR), Special transplanting technique (STP), Sustainable sugarcane initiating (SSI)technology.

Biotechnological method: Micropropagation

Micropropagation (Dinesh *et.al*,2015) is the best methodology available for rapid multiplication and production of quality seed free of any disease and pest which ensures maximum production or varieties(Urges *et.al.*,2021). Though this technique from one shoot apex about 180000 plants can be produced within a year which can cover about 14 hactares of nursery area.

Protocol: Multiple shoots were observed on Murashige and Skoog’s medium (MS medium) supplemented with BAP 0.2 mg/l and kinetin 0.1 mg/l. Best rooting was observed on MS media supplemented with IAA 1 mg/l and NAA 1 mg/l (table1).Soil mixture containing (soil+vermiculite+sand in 4:1:1 proportion by volume). 2 ft distance between 2 plants and 3 ft distance between 2 rows was found as an appropriate method for planting tissue cultured plantelets in the field. Setts obtained from tissue culture plants planted at 15 cm spacing gave higher yield than the other spacing.



Table1. Media composition

Ingredients	Shoot apex medium (mg/l) (whites basal)	Multiplication medium (mg/l) (modified MS)	Rooting medium (mg/l) (whites medium)
Ammonium nitrate	-	1640	-
Potassium nitrate	-	1900	-
Calcium chloride	-	440	-
Magnesium sulphate	720	-	720
Sodium sulphate	200	-	200
Potassium chloride	65	-	65
Potassium nitrite	80	-	80
Calcium nitrite	300	-	300
Sodium dihydrogen orthophosphate	16.5	-	16.5
Potassium dihydrogen orthophosphate	-	170	-
Boric acid	1.5	6.2	1.5
Cobalt chloride	-	0.025	-
Sodium molybdate	-	0.25	-
Cupric sulphate	-	0.25	-
Potassium iodide	0.75	-	0.75
Zinc sulphate	3	-	3
Manganese sulphate	7	22.5	7
Ferric EDTA	36.7	36.7	36.7
Meso inositol	100	100	-
Kinetin	1.07	1.07	-
Gibberlic acid	0.5	0.5	-
Indole Butyric acid	1	-	-
Glycine	2	2	3
Nicotinic acid	-	0.5	0.5
Pyridoxine	-	0.5	0.1
Thiamine	-	0.1	0.1
Naphthalene acetic acid	-	0.5	1
6 Benzyl amino purine	-	0.25	-
Calcium pantithenate	-	-	1
Riboflavin	-	-	1
Coconut water	100 ml	100 ml	-
Sucrose	20 g	20 g	20 g
pH	5.8	5.8	7.0

ENGINEERING RESEARCH

Micropropagation chiefly consist of four major steps:

(a) Collection of explant and sterilization : shoots(Purnamaningsih.*et.al*,2021)are collected from 3-4 months old crop and selection of apices of 10 cm long ,outer sheaths are removed by rectified spirit. Sample thoroughly rinsed in 70% ethanol for 1 minute. Disinfection is done by treating with chlorine water or sodium hypochloride solution for 10-15 minutes. The smell of chlorine is removed by 3-4 washing of sterile water under aseptic condition.

(b) Initial explant culture:

Apex meristem put in laminar flow where explant(Purnamaningsih *et.al.*,2021) placed aseptically on MS medium.

The basal MS medium along with suitable conc.of auxin and cytokinin is used for multiplication (Patel *et.al.*,2018)

(C) Multiplication:

2 to 5 shoots in 1st multiplication cycle(Baday *et.al.*,2020)of about 45 days.It is advised not to go beyond 7th cycles of sub-culturing because after 7th cycle,a green mass start to appear at base of formed shoot. The number of resulting shoots under favourable conditions may produce 36000 to 75000 plants depending on the genotypes.


(d)Rooting :

It achieved by transferring the individual or group of plants in rooting medium , root initiation is visible in many genotypes in 1 to 3 weeks . During tissue culture(Hendre *et.al.*,1983)*adventitious* roots or shoots can be induced by transferring the callus to medium containing different ratios of auxin and cytokinin.

(e) Hardening:It means transfer of rooting plantelets in field or pots which is also known as acclimatisation. Damaged plants should be eliminated. A mixture of sterilized sieved soil and sand in the ratio of 2:1 should be used for transplanting. The hardening process takes about 20-30 days. The hardened plant is transferred in the field in trenches at a distance of 45 cm or 60 cm within row and 90 cm between rows . Intercultural practices in crop raised through tissue culture are similar to conventional method.

Merit: There are certain merits of micropropagation(Sawant *et.al.*,2014) that's why it attained a certain degree of popularity in field of single bud propagation through tissue culture or biotechnological technique. Merits of micropropagation are: quick multiplication(Hendre *et.al.*,1983), disease free material ,true to type plants, easier transport, rejuvenation of old varieties,germplasm storage,micropropagated plants are more vigorous, give higher cane yield and sucrose %. The quality of seed produced by this technique can be maintained for 3 to 4 years.

fig 2 : Different steps of micropropagation in sugarcane



Micropropagation of sugarcane-laboratory. (a) Actively growing top. (b) Spindle. (c) Establishment of shoot cultures in vitro. (d) Rapid shoot multiplication through subculturing. (e) Rate of shoot multiplication per cycle. (f) Root formation in shoots. (g) Hardening of plantlets ex vitro

Conventional methods:

Material required: single bud setts, sett cutting machine, solution of malathian and bavistin 2%, gunny bags, cuvettes, polytrays, micro irrigation system, NPK ratio (19:19:19), specially designed racks.

Method: 1: Single budded nodes obtained with the help of sett cutting machine.

2. Nodes with sound buds are filled in gunny bags and dipped in solution of malathian and bavistin 2%

3. These treated nodes in gunny bags are kept under shade for six days to allow the buds to sprout and root initiation.

4. Water curing every day in morning and evening times

5. Workers are filling the cuvettes with root media.

6. Sprouting nodes are transferred to the cuvettes or holes of polytrays.

7. Unsprouted/ unrooted nodes are rejected in the second grading.
8. Transfer of polytrays in nursery farm.
9. Microirrigation system in nursery farm.
10. Sprouting nodes are coming outside.
11. N:P:K is 19:19:19 solution is sprayed on the growing seedlings at 10 days interval for vigorous growth.
12. Polytrays which are half filled with the rooting media ,followed by complete filling, pressing/compacting with fingers.
13. Workers are doing in grading for ready to issue the farmers.
14. Seedlings are ready in our nursery farm to issue the farmers.
15. Ready to issue the disease free sugarcane seedlings.
16. Healthy plants are ready to issue for farmers.
17. Sugarcane seedlings are transferred from nursery farm to sugarcane growers by tracks which are specially manufactured.

Merit: Single bud propagation completely eliminates the impact of the apical dominance, therefore emergence of neat, healthy and sturdy tillers .

It promotes uniform growth of sugarcane by improving luminous energy and fertilizers. Effort come under the sustainable sugarcane initiative (SSI) which aim to focus on using less seeds with wider seed spacing and better water and nutrient management to increase the cane yield significantly.

It has been observed that due to saving in seed cane material, the highest net return can be obtained with setting raised from single bud setts. According to study about 80% of seed material can be saved by planting single bud setts over the conventional 3 bud setts planting . The food reserves and moisture content in the single bud setts deplete at a faster rate compared to 2 or 3 bud setts, therefore sufficient soil moisture and nutrient are pre-requisite for viability of single bud setts as planting material.In view of the benefits of using single bud setts for sugarcane planting ,extensive field experiments were carried out at ICAR- IISR, Lucknow and its validation was also done at the farmers field in different sugar factory zones of Uttar Pradesh.

Mission: To fulfill the requirement of sugarcane with economical plantation. To revolutionize the field of sugarcane production through research and development with integrated viable technologies. To be a vibrant force in the sugarcane based industries and facilitate sustainable prosperity for all its participant.

Vision: The culmination of an enterprises endorsed with technological reference for sugarcane based industries.

A dedicated approach to always deliver high yield and disease free sugarcane.

Update: Development of protocol for tissue culture of sugarcane and development of agronomy for tissue cultured plants was essential to make use of tissue culture(Baday *et.al.*,2020) plantelets in three tier system of sugarcane seed production. The nutritional requirements for in vitro propagation protocol of sugarcane is highly dependent on the genotype and explant used .In vitro propagation that enables the rapid and large scale production of disease free planting material as being exercised with different crops is a prerequisite .

Result indicated that 15 days interval is the best optimum period of subculturing to enhance the rate of multiplication and produce more number of elongated and healthy shoots in a seed multiplication program through in vitro micropropagation

Future prospects: sugar is becoming an essential commodity to fulfill the requirement of ever increasing demand of population especially in contrast with South Asian and African continent. Our chief goal will be maximize sugarcane production in less amount of seed sett. Biotechnological route in form of micropropagation assured guarantee to fulfill this goal at rapid pace in comparison with conventional methods. Construct of tissue culture laboratory should be mandatory in every sugarcane industry unit to encourage rapid multiplication of single bud propagation.

Reference:

- (1)Baday,S.J. Sugarcane of rapid multiplication by calligenesis.JPSC 2020 1660,012005
- (2)D.Patel,M.P.Chaudhary, Influence of plant growth regulators on growth, yield and quality of sugarcane under South Gujarat,J.Pharmacogn. Phytochem.7(IS)(2018)9597.
- (3)Dinesh,P.,Saraniya,A.R., Ramanathan,T.In vitro studies of sugarcane variety co-91017 through micropropagation of shoot tip culture.Adv.plants Agric.Res.2015,2,234-238.
- (4)G.D.Urgesa,E.O.Keyatta, Effect of harvesting ages on yield component of sugarcane varieties, Ethiopia,Int J Food sci 2021(2021)
- (6)Hendre R R Iyer RS,Kotwal M ,Khupse SS,(1983) Rapid multiplication of sugarcane by tissue culture.sugarcane 1:4-9
- (7)Salokhe,S. Development of an efficient protocol for tissue culture of sugarcane.Plant cell Biotech.Mol.Bio,2021,4,9-21.
- (8)Sawant RA,Tawar PN,Meti NT, Ranjekar PK . Role of sugarcane micropropagation for production of quality seed. Int. Journal of recent Biotechnology.2014;2:34-41
- (9)Purnamaningsih R; Sukmadjaja,D; Suhesti,S., Raghu,S. In vitro propagation of six selected sugarcane mutant clones through leaf explants IOP conf.ser. Earth Environment.sci,2021,883,012075.